











**MODIFIED AMINOACIDS, PHARMACEUTICALS CONTAINING THESE COMPOUNDS AND METHOD FOR THEIR PRODUCTION****Publication number:** WO9811128**Publication date:** 1998-03-19**Inventor:** RUDOLF KLAUS (DE); EBERLEIN WOLFGANG (DE); ENGEL WOLFHARD (DE); PIEPER HELMUT (DE); DOODS HENRI (DE); HALLERMAYER GERHARD (DE); ENTZEROOTH MICHAEL (DE); WIENEN WOLFGANG (DE)**Applicant:** THOMAE GMBH DR K (DE); RUDOLF KLAUS (DE); EBERLEIN WOLFGANG (DE); ENGEL WOLFHARD (DE); PIEPER HELMUT (DE); DOODS HENRI (DE); HALLERMAYER GERHARD (DE); ENTZEROOTH MICHAEL (DE); WIENEN WOLFGANG (DE)**Classification:****- international:** C07K5/06; C07K5/065; G01N33/94; A61K38/00; C07K5/00; G01N33/94; A61K38/00; (IPC1-7): C07K5/06; C07D401/12; C07D409/12; C07D417/12; C07D419/12**- european:** C07K5/06A2; C07K5/06T; G01N33/94B**Application number:** WO1997EP04862 19970908**Priority number(s):** DE19961036623 19960910; DE19971020011 19970514**Also published as:** EP0927192 (A1)  
 US6344449 (B1)  
 JP2003300959 (A5)  
 EP0927192 (A0)  
 BR9712023 (A)  
 EP0927192 (B1)  
 TR9900537T (T2)  
 BG64214 (B1)  
 AU721035C (C)  
 AU721035B (B2)

less &lt;&lt;

**Cited documents:** WO9615148  
 EP0706999  
 US4826870  
 US4873342  
 US5310743  
 EP0415413  
 EP0284632  
 EP0298135  
 WO9604928  
 EP0723774

less &lt;&lt;

**Report a data error here****Abstract of WO9811128**

The invention concerns modified aminoacids of general formula (I) in which A, Z, X, n, m, R, R<2>, R<3>, R<4> and R<11> are defined in claims 1 through 5, their tautomers, their diastereomers, their enantiomers, their mixtures and their salts, specially their physiologically compatible salts with inorganic or organic acids or bases, pharmaceuticals containing these compounds, their use and the method for their production, as well as their use for the production and purification of antibodies and as marked compounds in RIA and ELISA assays and as diagnostic or analytic auxiliary agents in neurotransmitter research.

---

Data supplied from the **esp@cenet** database - Worldwide

**THIS PAGE BLANK (USPTO)**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K</b>		A2	(11) International Publication Number: <b>WO 98/05292</b>
			(43) International Publication Date: 12 February 1998 (12.02.98)
(21) International Application Number: PCT/US97/13383 (22) International Filing Date: 6 August 1997 (06.08.97) (30) Priority Data: 08/700,628                      8 August 1996 (08.08.96)                      US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (72) Inventors: LOWE, Derek, B.; 237 North 9th Street, Kenilworth, NJ 07033 (US). CHANG, Wei, K.; 63 West Cedar Street, Livingston, NJ 07039 (US). KOZLOWSKI, Joseph, A.; P.O. Box 7391, Princeton, NJ 08543-7391 (US). BERGER, Joel, G.; 5675 North Camino Esplandora, Tucson, AZ 85718 (US). McQUADE, Robert; 221 Mountainview Avenue, Scotch Plains, NJ 07076 (US). BARNETT, Allen; 13 Flanders Drive, Pine Brook, NJ 07058 (US). SHERLOCK, Margaret; 34 Parkway West, Bloomfield, NJ 07003 (US). TOM, Wing; 133 Cedar Grove Parkway, Cedar Grove, NJ 07009 (US). DUGAR, Sundee; 749 Wingate Drive, Bridgewater, NJ 08807 (US). CHEN, Lian-Yong; 10 Pond View Drive, Plainsboro, NJ 08536 (US). CLADER, John, W.; 428 North Union Avenue, Cranford, NJ 07016 (US). CHACKALAMANNIL, Samuel; 79 Stratford Road, East Brunswick, NJ 08816 (US). WANG,			Yuguang; 38 Princess Drive, North Brunswick, NJ 08902 (US). McCOMBIE, Stuart, W.; 28 Hanford Place, Caldwell, NJ 07006 (US). TAGAT, Jayaram, R.; 133 Boynton Court, Westfield, NJ 07090 (US). VICE, Susan, F.; 1144 Sawmill Road, Mountainside, NJ 07092 (US). VACCARO, Wayne, D.; 1706 Westover Road, Yardley, PA 19067 (US). GREEN, Michael, J.; 1146 Rancho Encinitas Drive, Encinitas, CA 92024 (US). BROWNE, Margaret, E.; 26 Highfield Road, Bloomfield, NJ 07003 (US). ASBEROM, Theodros; 525 Eagle Rock Avenue, West Orange, NJ 07052 (US). BOYLE, Craig, D.; 180 Northfield Road, Bridgewater, NJ 08807 (US). JOSIEN, Hubert, B.; 5441 Washington Boulevard, Jersey City, NJ 07310 (US). (74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.
(54) Title: MUSCARINIC ANTAGONISTS			
<div style="text-align: right;">(I)</div>			
(57) Abstract			
Di-N-substituted piperazine or 1,4 di-substituted piperidine compounds in accordance with formula I (including all isomers, salts, esters, and solvates) wherein R, R <sup>1</sup> , R <sup>2</sup> , R <sup>3</sup> , R <sup>4</sup> , R <sup>21</sup> , R <sup>27</sup> , R <sup>28</sup> , X, Y and Z as defined herein are muscarinic antagonists useful for treating cognitive disorders such as Alzheimer's disease. Pharmaceutical compositions and methods of preparation are also disclosed. Also disclosed are synergistic combinations of compounds of the above formula with acetylcholinesterase inhibitors.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

10

15

## MUSCARINIC ANTAGONISTS

20

### BACKGROUND OF THE INVENTION

The present invention relates to di-N-substituted piperazines and 1,4-di-substituted piperidines useful in the treatment of cognitive disorders, pharmaceutical compositions containing the compounds, methods of treatment using the compounds, and to the use of said compounds in combination with acetylcholinesterase inhibitors.

Alzheimer's disease and other cognitive disorders have received much attention lately, yet treatments for these diseases have not been very successful. According to Melchiorre et al. (J. Med. Chem. (1993), 36, 3734-3737), compounds that selectively antagonize M2 muscarinic receptors, especially in relation to M1 muscarinic receptors, should possess activity against cognitive disorders. Baumgold et al. (Eur. J. of Pharmacol., 251, (1994) 315-317) disclose 3- $\alpha$ -chloroimperialine as a highly selective m2 muscarinic antagonist.


The present invention is predicated on the discovery of a class of di-N-substituted piperazines and 1,4-di-substituted piperidines, some of which have m2 selectivity even higher than that of 3- $\alpha$ -chloroimperialine. Logemann et al (Brit. J. Pharmacol. (1961), 17, 286-296) describe certain di-N-substituted piperazines, but these are different from the

-2-

inventive compounds of the present invention. Furthermore, the compounds of Logemann et al. are not disclosed to have activity against cognitive disorders.

## 5 SUMMARY OF THE INVENTION


The present invention relates to compounds according to the structural formula I,




including all isomers and pharmaceutically acceptable salts, esters, and solvates thereof,

wherein one of Y and Z is N and the other is N, CH, or C-alkyl;





X is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>6</sup>-, -CO-, -CH<sub>2</sub>-, -CS-, -C(OR<sup>5</sup>)<sub>2</sub>-, -C(SR<sup>5</sup>)<sub>2</sub>-, -CONR<sup>20</sup>-, -C(alkyl)<sub>2</sub>-, -C(H)(alkyl)-, -NR<sup>20</sup>-SO<sub>2</sub>-, -SO<sub>2</sub>-NR<sup>20</sup>-, -NR<sup>20</sup>CO-, -O-CO-NH-, -NH-CO-O-,










R is




-3-



5


(wherein X<sup>1</sup> is -CH<sub>2</sub>-, -O-, or -NR<sup>7</sup>-),

-4-



hydrogen, acyl, alkyl, alkoxy, alkenyl, cycloalkyl, cycloalkyl substituted with one or two groups selected from the group consisting of alkyl and carbonyl, cycloalkenyl, bicycloalkyl, arylalkenyl, benzyl, benzyl substituted with up to three independently selected R<sup>3</sup> groups, cycloalkylalkyl, polyhaloacyl, benzyloxyalkyl, hydroxyC<sub>2</sub>-C<sub>20</sub>alkyl, alkenylcarbonyl, alkylarylsulfonyl, alkoxycarbonylaminoacyl, alkylsulfonyl, or arylsulfonyl, additionally, when X is -CH<sub>2</sub>-, R may also be -OH; in further addition, when X is not N, R may also be hydroxymethyl, in further addition, R and X may combine to form the group Prot-(NOAA)<sub>r</sub>-NH- wherein r is an integer of 1 to 4, Prot is a nitrogen protecting group and when r is 1, NOAA is a naturally occurring amino acid or an enantiomer thereof, or when r is 2 to 4, each NOAA is a peptide of an independently selected naturally occurring amino acid or an enantiomer thereof;

R<sup>1</sup> and R<sup>21</sup> are independently selected from the group consisting of H, alkyl, alkenyl, cycloalkyl, cycloalkenyl, bicycloalkyl, alkynyl, cyano, aminoalkyl, alkoxy carbonyl, aminocarbonyl, hydroxyamidino, alkoxy carbonylalkyl, phenyl alkyl, alkylcarbonyloxyalkyl,






-5-

H, —OH, (provided  $R^1$  and  $R^{21}$  are both not —OH and Y is not N), formyl, —CO alkyl, —COacyl, —COaryl, and hydroxyalkyl; additionally  $R^1$  and  $R^{21}$  together may form the group  $=CH_2$ ,  $=N-OR^5$ ,  $=N-CN$ ,  $=N-N(R^5)_2$ ,


5  $=CH-alkyl$ , alkylene,  $=C-alkyl$  or  $=C(halo)_2$ ; in further addition,  $R^1$  and  $R^{21}$  together with the carbon atom to which they are attached may



or  $R^1$  and  $R^{21}$  together with the carbon atom to which they are attached may form a saturated heterocyclic ring containing 3 to 7 carbon atoms, one or more of which may be optionally substituted by alkyl, and one or

10 two groups independently selected from S, O, and N- $R^{20}$ ;

$R^2$  is



$R^3$ ,  $R^4$ ,  $R^{22}$ ,  $R^{24}$ , and  $R^{25}$  are independently selected from the group consisting of alkyl, H, halo, alkoxy, benzyloxy, benzyloxy substituted by nitro or aminoalkyl, haloalkyl, polyhaloalkyl, nitro, cyano,

15 sulfonyl, hydroxy, amino, alkylamino, formyl, alkylthio, polyhaloalkoxy, acyloxy, trialkylsilyl, alkylsulfonyl, arylsulfonyl, acyl, alkoxycarbonyl alkylsulfinyl;  $-OCONH_2$ ,  $-OCONH-alkyl$ , alkylaminoalkyl, dialkylaminoalkyl,  $-COOH$ ,  $-CON(R^{20})_2$ ,  $-OCON(alkyl)_2$ ,  $-NHCOO-alkyl$ ,

20  $-NHCO-alkyl$ , phenyl, hydroxyalkyl, or morpholino;

each  $R^5$  and  $R^6$  is independently selected from the group consisting of H and alkyl, provided that when X is  $C(OR^5)_2$  or  $C(SR^5)_2$ , both  $R^5$  groups cannot be H, and in addition, when X is  $C(OR^5)_2$  or  $C(SR^5)_2$ , the two  $R^5$  groups in X may be joined to form  $-(CR^{20}_2)_p$

25 wherein p is an integer of 2 to 4;

$R^7$  is independently selected from the group consisting of H, alkyl, arylalkyl, cycloalkyl, aryl and aryl substituted with  $R^3$  and  $R^4$  as defined herein;

-6-

each  $R^8$  is independently selected from the group consisting of H, hydroxyalkyl or alkyl, or two  $R^8$  groups may be joined to form an alkylene group;

$R^9$  is H, alkyl, aralkyl, or acyl;

5  $R^{20}$  is H, aryl or alkyl;

$R^{27}$  and  $R^{28}$  are independently selected from the group consisting of H, alkyl, hydroxyalkyl, arylalkyl, aminoalkyl, haloalkyl, thioalkyl, alkylthioalkyl, carboxyalkyl, imidazolylalkyl, and indolylalkyl; or  $R^{27}$  and  $R^{28}$  may combine to form an alkylene group;

10  $R^{29}$  is H, alkyl, -CO-alkyl, -CO-cycloalkyl, alkoxy carbonyl, amino carbonyl, aryloxy carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfonyl, arylsulfonyl or -SO<sub>2</sub>-NH- $R^{20}$ ;

$R^{30}$  is H, alkyl, aryl, cycloalkyl, hydroxyalkyl, aminoalkyl, -COOR<sup>20</sup>, -CON( $R^{20}$ )<sub>2</sub> or cyano;

15  $R^{31}$  and  $R^{32}$  are the same as  $R^{30}$  and in addition, two  $R^{30}$ ,  $R^{31}$  and  $R^{32}$  groups may form the group -(CH<sub>2</sub>)<sub>r</sub> (wherein r is 1 to 6), in further addition,  $R^{31}$  and  $R^{32}$  can also be hydroxy, -N( $R^{20}$ )<sub>2</sub>, -O-acyl, -N( $R^{20}$ )acyl, -OCOOR<sup>20</sup>, or -OCON( $R^{20}$ )<sub>2</sub>;

20  $R^{33}$  is aryl or heteroaryl, with the proviso that when  $R^{33}$  is heteroaryl, the CO- $R^{33}$  bond is to a carbon atom in the  $R^{33}$  group;


$R^{34}$  is alkyl, cycloalkyl or aryl and in addition  $R^{34}$  may also be H when  $R^1$  and  $R^{21}$  together with the carbon atom to which they are attached form a saturated heterocyclic ring containing 3 to 7 carbon atoms and two groups independently selected from S, O, and N- $R^{20}$ ;

25  $R^{35}$  is -CH<sub>2</sub>-, -NR<sup>20</sup>- or -O-;

$R^{36}$  is -NH<sub>2</sub>, alkyl or alkoxy;

$R^{37}$  is independently selected from the group consisting of H and alkyl;

$R^{38}$  is -CO-(CH<sub>2</sub>)<sub>0-5</sub>-OR<sup>5</sup>, -SO<sub>2</sub>-(alkyl), or



30

wherein  $q_1$  and  $q_2$  are independently 1-5, provided that the sum of  $q_1$  and  $q_2$  is 2-5; and


-7-

$R^{39}$  and  $R^{40}$  are independently selected from the group consisting of  $=O$  and  $(H,H)$ .

In a preferred group of compounds  $Y$  and  $Z$  are  $N$ .

In another preferred group of compounds  $Y$  is  $CH$  and  $Z$  is  $N$ .


5 In another preferred group of compounds  $R$  is



and  $X$  is  $O$ ,  $SO$ ,  $SO_2$ ,  $CH_2$ ,  $CH(alkyl)$ ,  $C(alkyl)_2$ ,  $-CH(OH)-$ , or  $-N(R^{20})CO$ .

10 In another preferred group of compounds  $R^3$  and  $R^4$  are  $H$  and either  $R^1$  is cycloalkyl, or alkyl, and  $R^{21}$  is  $H$  or  $R^1$  and  $R^{21}$  together form  $=O$ .

In another preferred group of compounds  $R$  is




$X$  is  $O$ ,  $SO$ ,  $SO_2$ ,  $CH_2$ ,  $CH(alkyl)$ ,  $C(alkyl)_2$ , or  $-N(R^{20})CO$ ;  $R^3$  and  $R^4$  are  $H$  and either  $R^1$  is cycloalkyl or alkyl; and  $R^{21}$  is  $H$  or  $R^1$  and  $R^{21}$

15 together form  $=O$  or

In another preferred group of compounds at least one of  $R^{27}$  and  $R^{28}$  is alkyl.

In another preferred group of compound one of  $R^{27}$  or  $R^{28}$  is methyl and the other is hydrogen.


20 In another preferred group of compounds  $R$  is




In another preferred group of compounds  $X$  is  $SO_2$ ,  $CH_2$ , or  $-N(CH_3)-CO-$ .

In another group of preferred compounds,  $R^2$  has the formula


-8-



and  $R^{30}$  is H or  $CH_3$ ;  $R^{31}$  and  $R^{32}$  are H; and  $R^{33}$  is ortho-substituted aryl or heteroaryl, preferably



5 In another group of preferred compounds,  $R^2$  has the formula



$R^{34}$  is methyl and  $R^{31}$  and  $R^{32}$  are H.

Preferred specific compounds of this invention are listed

hereinafter as compound numbers: 17, 18, 25, 30, 31, 32, 34, 35,  
 10 36, 37, 41, 43, 44, 49, 53, 54, 56, 57, 58, 59, 80, 82, 84, 85,  
94, 98, 100, 108, 121, 126, 127, 137, 145, 151, 152, 154,  
155, 162, 166, 178, 179, 181, 185, 190, 191, 194, 199, 214,  
215, 216, 225, 247, 253, 256, 257, 337, 339, 340, 341, 349,  
351, 367, 409, 459, 479, 488, 489, 490, 500, 501, 502, 503,  
 15 505, 506, 507, 515, 516, 517, 555, 562.

Another aspect of the invention is a pharmaceutical composition which comprises a compound having structural formula I as defined above in combination with a pharmaceutically acceptable carrier.

20 Another aspect of the invention is the use of a compound formula I for the preparation of a pharmaceutical composition useful in the treatment of cognitive disorders and neurodegenerative diseases such as Alzheimer's disease.

Yet another aspect of the invention comprises a method for making a pharmaceutical composition comprising mixing a compound of  
 25 formula I with a pharmaceutically acceptable carrier.

-9-

Another aspect of this invention is a method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a compound of formula I.

5 Another aspect of this invention is a method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a combination of a compound of formula I with an acetylcholinesterase inhibitor.

10 Another aspect of this invention is a kit comprising in separate containers in a single package pharmaceutical compounds for use in combination to treat cognitive disorders in one container a compound of formula I in a pharmaceutically acceptable carrier and in a second container an acetylcholinesterase inhibitor in a pharmaceutically acceptable carrier, the combined quantities being an effective amount.

15

#### DETAILED DESCRIPTION

Except where stated otherwise the following definitions apply throughout the present specification and claims. These definitions apply regardless of whether a term is used by itself or in combination with  
20 other terms. Hence the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portions of "alkoxy", "haloalkyl", etc.

Alkyl represents a straight or branched saturated hydrocarbon chain having 1 to 20 carbon atoms, more preferably 1 to 8 carbon atoms.

25 Alkenyl represents a straight or branched hydrocarbon chain of from 2 to 15 carbon atoms, more preferably 2 to 12 carbon atoms, having at least one carbon-to-carbon double bond.

Alkynyl represents a straight or branched hydrocarbon chain of from 2 to 10 carbon atoms, more preferably 2 to 8 carbon atoms, having at least one carbon-to-carbon triple bond.

30 Cycloalkyl represents a saturated carbocyclic ring having 3 to 12 carbon atoms.

Cycloalkenyl represents a carbocyclic ring having from 5 to 8 carbon atoms and at least one carbon-to-carbon double bond in the ring.

35 Bicycloalkyl represents a saturated bridged carbocyclic ring having 5 to 12 carbon atoms.

-10-

Acyl represents a radical of a carboxylic acid having the formula alkyl-CO-, aryl-CO-, aralkyl-CO-, cycloalkyl-CO-, alkylcycloalkyl-CO-, and heteroaryl-CO-.


Halo represents fluoro, chloro, bromo or iodo.

5 Aryl represents phenyl or naphthyl optionally substituted with 1 to 5 R<sup>3</sup> groups.

Heteroaryl represents a cyclic group of 5 or 6 atoms, or a bicyclic group of 9 or 10 atoms, at least one of which is carbon and having at least one O, S, or N atom interrupting a carbocyclic ring having a  
10 sufficient number of pi electrons to provide aromatic character. Carbon atoms may optionally be substituted by R<sup>3</sup> groups. Nitrogen atoms may optionally be substituted -R<sup>20</sup> or -COR<sup>20</sup> groups. Preferred heteroaromatic groups are pyridyl, furanyl, benzofuranyl, thienyl, benzothienyl, thiazolyl, thiadiazolyl, imidazolyl, pyrimidinyl, pyrazinyl,  
15 pyridazinyl, 1,3,5-triazinyl, and indolyl.

Polyhalo represents substitution of at least 2 halo atoms to the group modified by the term "polyhalo".

Hydroxyamidino represents a group having the formula



20 Azabicyclo represents a saturated bridged ring containing from 4 to 8 carbon atoms and at least one nitrogen atom.

Sulfonyl represents a group of the formula -SO<sub>2</sub>-.

Sulfinyl represents a group of the formula -SO-.

25 Alkylene represents a group having the formula -(CH<sub>2</sub>)<sub>q</sub>, wherein q is an integer of from 1 to 20.

Naturally occurring amino acid (NOAA) means an acid selected from the group consisting of alanine (ala), arginine (arg), asparagine (asn), aspartic acid (asp), cysteine (cys), glutamine (gln), glutamic acid  
30 (glu), glycine (gly), histidine (his), isoleucine (ile), leucine (leu), lysine (lys), methionine (met), phenylalanine (phe), proline (pro), serine (ser), threonine (thr), tryptophan (trp), tyrosine (tyr), and valine (val).

Nitrogen protecting group (Prot) means a group capable of protecting a nitrogen on a naturally occurring amino acid (or an  
35 enantiomer thereof) from reaction. Preferred nitrogen protecting groups

-11-

are carbobenzyloxy (CBZ),  $\text{CH}_3\text{OCO}(\text{CH}_2)_9\text{CO}$ , and t-butoxycarbonyl (BOC). Of course any operable nitrogen protecting group is included.

When a variable appears more than once in the structural formula, for example  $\text{R}^5$  when X is  $-\text{C}(\text{OR}^5)_2-$ , the identity of each  
5 variable appearing more than once may be independently selected from the definition for that variable.


Compounds of this invention may exist in at least two stereo configurations based on the asymmetric carbon to which  $\text{R}^1$  is attached, provided that  $\text{R}^1$  and  $\text{R}^{21}$  are not identical. Further stereoisomerism is  
10 present when X is SO, or  $\text{C}(\text{OR}^5)_2$  (when the two  $\text{R}^5$  groups are not the same) or when  $\text{R}^{27}$  or  $\text{R}^{28}$  is not hydrogen. Also within formula I there are numerous other possibilities for stereoisomerism. All possible stereoisomers of formula I are within the scope of the invention.

Compound of formula I can exist in unsolvated as well as solvated  
15 forms, including hydrated forms. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol and the like, are equivalent to the unsolvated forms for purposes of this invention.

A compound of formula I may form pharmaceutically acceptable salts  
20 with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base forms with a sufficient  
25 amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia or sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in  
30 certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base forms for purposes of the invention.




Compounds in accordance with formula I may be produced by  
processes known to those skilled in the art as shown by the following  
35 reaction steps:

-12-

**General Description of Methods:**Compounds wherein R<sup>2</sup> has the formula

can be made as shown in Scheme I.

5

**Scheme 1**For compounds wherein R<sup>2</sup> has the formula




-13-


- Intermediate III is treated with a suitably activated carboxylic acid derivative,  $R_{33}CO-X$ , where X is a leaving group such as halogen or  $OCOCH_3$ . Alternatively, a mixture of compound III and  $R_{33}COOH$  can be treated with N-hydroxybenzotriazole and a carbodiimide such as N-ethyl-N-(3-dimethylaminopropyl)-carbodiimide hydrochloride or dicyclohexylcarbodiimide in the presence of a base such as triethylamine and a solvent such as DMF to give I. Alternatively, intermediate IIIc is treated with a group  $R_{29}-X$ , where X and  $R_{29}$  are as previously defined (except  $R_{29} \neq H$ ), in the presence of a base such as triethylamine.

Intermediates III can be made via a variety of methods. When  $X=CH_2$  or  $CO$ , III can be made as shown in Scheme 2:

Scheme 2




-14-




- 5 Aryl halide **IV** is treated with an alkyllithium such as *n*-butyllithium or *t*-butyllithium followed by reaction with an  $\text{RCON}(\text{CH}_3)\text{OCH}_3$  to give compound **IIIa**, which can be deprotected and converted to **I** as shown above. Additionally, **IIIa** can be reduced to the alcohol **V** with a suitable reducing agent such as sodium borohydride. Compound **V** can also be made by treatment of **IV** with an alkyllithium as previously described followed by reaction with  $\text{RCHO}$ . Compound **V** is converted to
- 10 compounds **IIIb** by treating **V** with a reducing agent such as triethylsilane in the presence of a strong acid such as trifluoroacetic acid.

Compounds of formula **Ia**, **Ib**, and **Ic** can be prepared via a related sequence as shown in Scheme 3:

Scheme 3




-15-




Intermediate IIIa is prepared as shown in Scheme 2, deprotected and then converted to Ia using methods of Scheme 1. This can then be converted to Ib and Ic using methods of Scheme 2.

- 5 Other compounds of this invention can be prepared by methods similar to those described in Schemes 2-3 wherein the aryl lithium reagent derived from IV is reacted with various electrophiles. For instance, compounds where X is NHCO or N(alkyl)CO can be prepared as shown in Scheme 4:

Scheme 4



-16-




- Intermediate IV is reacted with an alkyllithium as previously described followed by addition of an isocyanate RNCO. The intermediate is deprotected and converted to compounds of type 1d as described in Scheme 1. Compounds of type 1d are converted to compounds 1e by reacting with an alkyl iodide such as methyl iodide in the presence of a suitable base such as sodium hydride.

Intermediate IV can be prepared via one of the following procedures:


10

Scheme 5

Intermediate IVa:




-17-



Benzylamine V is protected as its trifluoroacetamide, treated with a halogenating agent such as bromine, dibromodimethylhydantoin, or bis(trifluoroacetoxy)iodobenzene, and deprotected with aqueous base to give VI. This is treated with a carboxylic ester derivative containing a leaving group such as trifluoromethanesulfonate in the 2-position followed by chloroacetyl chloride to give VII. Treatment of VII with ammonia followed by reduction with  $\text{NaBH}_4/\text{BF}_3$  etherate gives VIII. Treatment of VIII with an N-BOC 4-piperidinone derivative preferable in the presence of a Lewis acid such as titanium tetraisopropoxide followed by a reducing agent such as sodium cyanoborohydride affords IVa.

### Scheme 6

Intermediate IVb:




-18-

A piperidine 4-carboxylic acid derivative IX is protected on nitrogen using trifluoroacetic anhydride and converted to the corresponding acid chloride XI using thionyl chloride. Intermediate XI is reacted with an arylhalide in the presence of a Lewis acid such as aluminum chloride to give XII. The carbonyl group of XII is protected by treatment with ethylene glycol in the presence of a strong acid such as toluenesulfonic acid. The nitrogen is deprotected using aqueous alcoholic base, and the resulting compound treated with an N-BOC 4-piperidinone derivative as described in Scheme 5 to afford IVb.

Compounds of formula If (where  $R_{30} \neq H$ ) and II are prepared as shown in Scheme 7:

Scheme 7



Intermediate XIII is treated with an N-BOC 4-piperidinone derivative preferably in the presence of a Lewis acid such as titanium


-19-

tetraisopropoxide followed by treatment with diethylaluminum cyanide to give XIV. This is treated with Grignard reagent  $R_{34}MgCl$  followed by hydrolysis with aqueous acid to give IIIc. Compound IIIc can be converted to compounds of type If and II using the method shown in

5 Scheme 1.


In addition to methods described above, compounds of formula Ig (where A is O, S or N- $R^{20}$ ) can be prepared as described in Scheme 8.

Scheme 8




-20-

In the compound  $\text{HA}-(\text{CH}_2)_{2-5}-\text{AH}$ , each A is independently O, S, or N- $\text{R}_{20}$ , and each  $\text{CH}_2$  group may optionally be substituted by one or more alkyl groups. Examples of the compound  $\text{HA}-(\text{CH}_2)_{2-5}-\text{AH}$ , include




- 5 Compound **XV** (where  $\text{R}''$  is either  $\text{R}_{30}$  or  $\text{R}_{34}$ ) is prepared as described in schemes 7 and 10. This is treated with  $\text{HA}-(\text{CH}_2)_{2-5}-\text{AH}$  optionally in the presence of an acid catalyst and a dehydrating agent. When A is O, the acid catalyst is preferably an organic protic acid such as toluenesulfonic acid and the dehydrating agent is triethyl orthoformate. When A is S, a Lewis acid such as boron trifluoride etherate serves as both acid catalyst and dehydrating agent. The resulting product **XVI** is hydrolyzed with a strong base such as potassium hydroxide to give **XVII**. This is converted to **1e** via
- 10 methods described in scheme 1.
- 15 When  $\text{X} = \text{S}, \text{SO},$  or  $\text{SO}_2$ , intermediates **III** are prepared as shown in the following schemes:

### Scheme 9






-21-



Scheme 10



- 5 The above reactions may be followed if necessary or desired by one or more of the following steps; (a) removing any protective groups

-22-

- from the compound so produced; (b) converting the compound so-produced to a pharmaceutically acceptable salt, ester and/or solvate; (c) converting a compound in accordance with formula I so produced to another compound in accordance with formula I, and (d) isolating a  
5 compound of formula I, including separating stereoisomers of formula I.

Based on the foregoing reaction sequence, those skilled in the art will be able to select starting materials needed to produce any compound in accordance with formula I.

- 10 In the above processes it is sometimes desirable and/or necessary to protect certain groups during the reactions. Conventional protecting groups, familiar to those skilled in the art, are operable. After the reaction or reactions, the protecting groups may be removed by standard procedures.

- 15 The compounds of formula I exhibit selective m<sub>2</sub> and/or m<sub>4</sub> muscarinic antagonizing activity, which has been correlated with pharmaceutical activity for treating cognitive disorders such as Alzheimers disease and senile dementia.

- 20 The compounds of formula I display pharmacological activity in test procedures designated to indicate m<sub>1</sub> and m<sub>2</sub> muscarinic antagonist activity. The compounds are non-toxic at pharmaceutically therapeutic doses. Following are descriptions of the test procedures.

#### MUSCARINIC BINDING ACTIVITY

- 25 The compound of interest is tested for its ability to inhibit binding to the cloned human m<sub>1</sub>, m<sub>2</sub>, m<sub>3</sub>, and m<sub>4</sub> muscarinic receptor subtypes. The sources of receptors in these studies were membranes from stably transfected CHO cell lines which were expressing each of the receptor subtypes. Following growth, the cells were pelleted and subsequently homogenized using a Polytron in 50 volumes cold 10 mM Na/K  
30 phosphate buffer, pH 7.4 (Buffer B). The homogenates were centrifuged at 40,000 x g for 20 minutes at 4°C. The resulting supernatants were discarded and the pellets were resuspended in Buffer B at a final concentration of 20 mg wet tissue/ml. These membranes were stored at -80°C until utilized in the binding assays described below.

-23-

Binding to the cloned human muscarinic receptors was performed using  $^3\text{H}$ -quinuclidinyl benzilate (QNB) (Watson et al., 1986). Briefly, membranes (approximately 8, 20, and 14  $\mu\text{g}$  of protein assay for the m1, m2, and m4 containing membranes, respectively) were incubated with  $^3\text{H}$ -QNB (final concentration of 100-200 pM) and increasing concentrations of unlabeled drug in a final volume of 2 ml at 25°C for 90 minutes. Non-specific binding was assayed in the presence of 1  $\mu\text{M}$  atropine. The incubations were terminated by vacuum filtration over GF/B glass fiber filters using a Skatron filtration apparatus and the filters were washed with cold 10mM Na/K phosphate buffer, pH 7.4. Scintillation cocktail was added to the filters and the vials were incubated overnight. The bound radioligand was quantified in a liquid scintillation counter (50% efficiency). The resulting data were analyzed for  $\text{IC}_{50}$  values (i.e. the concentration of compound required to inhibit binding by 50%) using the EBDA computer program (McPherson, 1985). Affinity values ( $K_i$ ) were then determined using the following formula (Cheng and Prusoff, 1973);

$$K_i = \frac{\text{IC}_{50}}{1 + \frac{\text{concentration of radioligand}}{\text{affinity (K}_D\text{) of radioligand}}}$$

Hence, a lower value of  $K_i$  indicates greater binding affinity.

To determine the degree of selectivity of a compound for binding the m2 receptor, the  $K_i$  value for m1 receptors was divided by the  $K_i$  value for m2 receptors. A higher ratio indicates a greater selectivity for binding the m2 muscarinic receptor.

#### RESULTS OF THE TESTS

Compound	K <sub>i</sub> (nM)			
	No.	m1	m2	m4
	<u>17</u>	47.33	0.14	2.26
	<u>18</u>	48.37	0.11	0.77
	<u>25</u>	337.68	0.55	6.51
	<u>31</u>	308.95	0.63	12.10
	<u>41</u>	29.9	0.06	1.47
	<u>44</u>	36.79	0.11	0.76

-24-

<u>84</u>	12.32	0.04	0.06
<u>94</u>	28.99	0.02	0.76
<u>100</u>	6497	0.12	3.38
<u>554</u>	0.81	0.01	0.04
<u>590</u>	29.33	0.03	0.80

For the compounds appearing in the TABLE OF COMPOUNDS the following range of muscarinic antagonistic activity was observed

m2: 0.01 nM to 106 nM

m1: 0.24 nM to 1900 nM

5 m4: 0.02 nM to 705 nM

For preparing pharmaceutical compositions from the compounds of formula I, compounds capable of enhancing ACh release, and ACh'ase inhibitors, pharmaceutically acceptable, inert carriers are admixed with the active compounds. The pharmaceutically acceptable carriers may be either solid or liquid. Solid form preparations include 10 powders, tablets, dispersible granules, capsules, cachets and suppositories. A solid carrier can be one or more substances which may also act as dilutents, flavoring agents, solubilizers, lubricants, suspending agents, binders or tablet disintegrating agents; it may also 15 be an encapsulating material.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection.

Also included are solid form preparations which are intended to 20 be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions. These particular solid form preparations are most conveniently provided in unit dose form and as such are used to provide a single liquid dosage unit.

25 The invention also contemplates alternative delivery systems including, but not necessarily limited to, transdermal delivery. The transdermal compositions can take the form of creams, lotions and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

-25-

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active components. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation such as packeted tablets, capsules and powders in vials or ampules. The unit dosage form can also be a capsule, cachet or tablet itself, or it may be the appropriate number of any of these in a packaged form.

The quantity of active compound in a unit dose preparation may be varied or adjusted from 1 mg to 100 mg according to the particular application and the potency of the active ingredient and the intended treatment. This would correspond to a dose of about 0.001 to about 20 mg/kg which may be divided over 1 to 3 administrations per day. The composition may, if desired, also contain other therapeutic agents.

The dosages may be varied depending on the requirement of the patient, the severity of the condition being treating and the particular compound being employed. Determination of the proper dosage for a particular situation is within the skill of those in the medical art. For convenience, the total daily dosage may be divided and administered in portions throughout the day or by means providing continuous delivery. When a compound of formula I or a compound capable of enhancing ACh release is used in combination with an acetylcholinesterase inhibitor to treat cognitive disorders these two active components may be co-administered simultaneously or sequentially, or a single pharmaceutical composition comprising a compound of formula I or a compound capable of enhancing ACh release and an acetylcholinesterase inhibitor in a pharmaceutically acceptable carrier can be administered. The components of the combination can be administered individually or together in any conventional oral or parenteral dosage form such as capsule, tablet, powder, cachet, suspension, solution, suppository, nasal spray, etc. The dosage of the acetylcholinesterase inhibitor may range from 0.001 to 100 mg/kg body weight.


The invention disclosed herein is exemplified by the following preparation and examples which should not be construed to limit the

-26-

scope of the disclosure. Alternative mechanistic pathways and analogous structures may be apparent to those skilled in the art.


In the following examples intermediate numerals have normal type and are enclosed by parenthesis, e.g., (5). Product compounds are in bold type and underlined, e.g., 5

**Example 1: Synthesis of Compound No. 4**



An ether (650 ml) solution of the bromide (15.5 g, 0.07 mol) was cooled in a dry ice acetone bath where t-BuLi (100 ml, 1.7 M) was added dropwise. After stirring for 3h, the sulfur powder (4.9 g, 0.15 mol) was added and stirring continued for an additional hour. The temperature was warmed to room temperature and stirring continued for 16h. After quenching with water, ether (Et<sub>2</sub>O) was added and the organic phase was washed with 5% HCl, water, 10% Na<sub>2</sub>CO<sub>3</sub> and then brine. The solution was concentrated and purified by chromatography with 5% ethyl acetate (EtOAc) / hexane ( $R_f = 0.7$ / CH<sub>2</sub>Cl<sub>2</sub>: hexane 1:1) with 33 g yellow solid collected.


The sample was dissolved in THF (50 mL) and cooled in an ice water bath where it was added to a mixture of LAH (0.91g, 0.02 mol) and 20 mL THF. Stirring was continued for 10 minutes, then warmed to room temperature and stirred for 1h. The reaction was diluted with EtOAc and quenched with water. 10% HCl was added and the organic layer separated, washed with water and brine. After concentration and drying under vacuum, 6.6 g of a yellow oil was collected.




The thiol (19 g, 124 mmol) was dissolved in DMF (180 ml) and cooled in an ice bath where NaH (4.9 g, 60% in oil) was added in portions. Stirring was continued for 1h, then warmed to room temperature and stirred for an additional 1h. The 4-fluoroacetophenone (18.7 g, 136 mmol) was added dropwise and then stirred at 70°C for 3h. The reaction was concentrated on a rotovap, diluted with EtOAc, and washed with

-27-


water and brine. The solution was evaporated on a rotovap and the residue chromatographed on silica gel (20% EtOAc/hexane) to collect 24.5g (73%) of a pale yellow solid.



- 5 To a solution of the alcohol (10.1 g, 40 mmol) in toluene (100mL) was slowly added the trimethyl boroxine (3.3 g, 27 mmol). After 4h, the volatiles were removed by distillation (oil bath temp=140-150°C). Cooling to room temperature was followed by addition of toluene (50mL), and removal of the volatiles by distillation (repeat again). The
- 10 residue was placed under high vacuum and the flask heated to 150°C. After drying under vacuum overnight the off white solid was dissolved in THF (100ml) and used without further purification.




- The ketone (36 g, 133 mmol) was dissolved in THF (200 ml),
- 15 followed by addition of the catalyst (8.8 g in 80 ml THF).  $\text{BH}_3(\text{CH}_3)_2\text{S}$  (42 ml, 2mol/L in THF) was added dropwise, and the reaction stirred for 30min.  $\text{CH}_3\text{OH}$  (200mL) was added and the solution concentrated. The residue was dissolved in EtOAc and washed with water and brine. The solution was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give 40.3 g
- 20 of a pink oil.




- A  $\text{CH}_2\text{Cl}_2$  (600 ml) solution of the sulfide (40.3g, 147mmol) was cooled in an ice water bath where MCPBA (79.7 g, 70%) was added in portions. After stirring for 1h the temperature was warmed to room
- 25 temperature and stirred for 4h. After diluting with  $\text{CH}_2\text{Cl}_2$ , the reaction

-28-


was washed with 10%  $\text{Na}_2\text{CO}_3$ , water, and brine. The solution was dried over,  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to collect 42.5g of an off white solid.



- 5 A  $\text{CH}_2\text{Cl}_2$  (250 ml) solution of the alcohol (21.4 g, 70 mmol) and triethyl amine ( $\text{Et}_3\text{N}$ ) (19.5 ml) was cooled in an ice water bath where methanesulfonyl chloride (6.5 ml) were added dropwise. After 1h the reaction was diluted with  $\text{CH}_2\text{Cl}_2$  and washed with 2% HCl, water, saturated  $\text{NaHCO}_3$ , water, brine, and dried over  $\text{Na}_2\text{SO}_4$ . Filtration and  
10 concentration gave 27g of an oil which was used without further purification.



- A mixture of the mesylate (made from 21.4 g alcohol), piperazine (18 g, 0.07 mol), and 2,2,6,6-tetramethyl piperidine (11.8 g, 0.084 mol) was stirred in  $\text{CH}_3\text{CN}$  (200 ml) at reflux overnight. After cooling to room temperature the  $\text{CH}_3\text{CN}$  was removed on a roto vap and replaced with EtOAc. The solution was washed with 10%  $\text{Na}_2\text{CO}_3$ , and water. After  
15 concentration the residue was chromatographed on silica gel using 1:3 EtOAc/hexane. 21.9g (60%) of a gum was collected.




20




-29-

A mixture of the CBZ amine (10.3g, 19.6 mmol), 100 ml water, CH<sub>3</sub>OH (100 ml), and 100 ml conc. HCl was heated in an oil bath at 100°C for 5h. The mixture was cooled in an ice bath and 50% NaOH was added until the pH=9. EtOAc was added, followed by separation and washing with water. The organic solution was concentrated and the residue chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH 50:1, saturated with NH<sub>4</sub>OH solution). The product was collected in 86% as a heavy oil.




The amine (1.0 g, 2.5 mmol), N-BOC piperidinone (0.5 g, 2.5 mmol), and HOAc (0.15 ml) were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 ml). NaBH(OAc)<sub>3</sub> (0.55 g, 3.8 mmol) was added in several portions and the solution stirred overnight. CH<sub>2</sub>Cl<sub>2</sub> was added and washed with saturated NaHCO<sub>3</sub>, water and brine. The solution was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was chromatographed using EtOAc. A colorless gum (0.96g) was obtained in 65% yield.



The N-BOC compound (0.87 g, 1.5 mmol) was dissolved in EtOAc (20 ml), and 6N HCl (3.5 ml) was added with stirring. After 2h, CH<sub>2</sub>Cl<sub>2</sub> was added and the solution was washed with saturated NaHCO<sub>3</sub>, and


-30-

dried over  $\text{Na}_2\text{SO}_4$ . After filtration the solution was concentrated and dried under vacuum to give a white powder (0.6g, 85%).




- 5 Stir for 20h. at RT a solution of intermediate (1) (0.025 g), N-hydroxybenzotriazole (HOBT) (0.01 g), benzoic acid (0.0125 g),  $\text{Et}_3\text{N}$  (0.01 mL) and N-ethyl-N-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) (0.025 g) in DMF (0.3 ml). Dilute with EtOAc, wash with water and with  $\text{NaHCO}_3$  solution, dry over  $\text{MgSO}_4$  and filter through a small plug of silica gel, washing with EtOAc. Evaporate to obtain the
- 10 free base form of the title compound. Dissolve this in  $\text{CH}_2\text{Cl}_2$  (0.2 mL) and add to a solution of HCl in dioxane (4M; 0.1 mL) in  $\text{Et}_2\text{O}$  (1.5 mL). Stir for 5 min., centrifuge, wash by suspension and centrifugation 3x with  $\text{Et}_2\text{O}$  and dry at RT in a stream of  $\text{N}_2$ , then at high vacuum to obtain the dihydrochloride. Mp: 215-225°C, with decomposition; Mass spectrum:
- 15  $\text{MH}^+ = 574$ .

#### Example 2: Synthesis of Compound No. 17



-31-

- Stir for 20h. a mixture of intermediate (1) from example 1, (0.05 g),  $\text{CH}_2\text{Cl}_2$  (1.5 ml), o-toluoyl chloride (0.1 ml) and 1N aq. NaOH (2 ml). Separate the organic phase, dry and evaporate. Purify by preparative tlc, eluting with 5%  $\text{CH}_3\text{OH}-\text{CH}_2\text{Cl}_2$ . Extract the major band with 50%  $\text{CH}_3\text{OH}-\text{CH}_2\text{Cl}_2$  and evaporate. Precipitate, wash and dry the hydrochloride (yield = 0.065 g) in the manner described in the foregoing preparation. Mp: 182-190°C with decomposition; Mass spectrum:  $\text{MH}^+ = 588$ .


**Example 3: Synthesis of Compound No. 28**

- This synthesis can be carried out using either intermediate (3a) or (3b) produced as shown in examples 10 and 11. To a solution of (3b) (1.86 g) in dry THF (30 ml) at -70°C under  $\text{N}_2$  add n-BuLi (2.5 M in hexanes; 1.6 ml), stir for 10 min. and add a solution of N-piperonoyl-N,O-dimethyl-hydroxylamine (0.83 g) in THF (1 ml). Stir without cooling for 2h., add EtOAc, wash with water, dry ( $\text{MgSO}_4$ ) and evaporate.
- 15

-32-


- Chromatograph on silica gel with EtOAc-CH<sub>2</sub>Cl<sub>2</sub> to obtain the major component intermediate (4) as a thick oil (1.21 g). Stir a solution of (4) (0.4 g) in EtOAc (24 ml) and conc. HCl (4.5 ml) at RT for 3h. Dilute with EtOAc, and wash with excess 1N NaOH solution. Extract with EtOAc, wash with saturated brine, dry and evaporate to obtain the NH compound. Dissolve this in CH<sub>2</sub>Cl<sub>2</sub> (10 ml), and stir for 20h. at RT with o-toluoyl chloride (0.15 g) and NaOH (0.8 g) in H<sub>2</sub>O (10 ml). Extract with CH<sub>2</sub>Cl<sub>2</sub>, dry (MgSO<sub>4</sub>), evaporate and chromatograph on silica gel with 4% CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>. Evaporate the pure fractions to obtain the free base as a foam (0.33 g) Dissolve a small portion this in CH<sub>2</sub>Cl<sub>2</sub> and precipitate the hydrochloride salt as described in previous preparations. M.p.: 200-210°C, with decomposition; Mass spectrum: MH<sup>+</sup> = 554

**Example 4: Synthesis of Compound No. 30**



- 15 Stir a solution of the product of example 3 (0.31 g), ethanol (15 ml) and sodium borohydride (0.042 g) at RT for 3h. Evaporate, add water, extract with EtOAc, dry, evaporate and chromatograph on silica gel with 10% CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub> to obtain the product (0.28 g) as a white foam, a mixture of diastereoisomers. Mp: 85-95°C; Mass spectrum:
- 20 MH<sup>+</sup> = 556.

**Example 5: Synthesis of Compound No. 31**




Stir a solution of the product of example 4 (0.1 g) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) and add triethylsilane (0.3 mL) and trifluoroacetic acid (0.12 mL). Stir at

-33-

RT for 72h. Add excess 1N NaOH solution, extract with EtOAc, dry, evaporate and purify by preparative tlc on silica plates, eluting with 5% CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>. Remove and extract the major band, evaporate the eluate, and convert to the hydrochloride in the usual manner, to obtain a  
 5 white powder (0.08 g). Mp: 210-215°C, with decomposition; Mass spectrum: MH<sup>+</sup> = 540.

**Example 6: Preparation of Compound No. 24**



To a solution of (3b) (Example 10) (1.38 g) in dry THF (20 ml) at  
 10 -70°C under dry N<sub>2</sub> add n-BuLi (2.5M in hexanes; 1.7 ml). Stir for 5 min.,  
 and add a solution of 3,4-methylenedioxyphenyl isocyanate (crude  
 product, prepared by heating the corresponding acyl azide in toluene)  
 (0.68 g) in THF (5 ml). Stir for 30 min without cooling, and work up in  
 EtOAc-aq.NaHCO<sub>3</sub>, dry (MgSO<sub>4</sub>) and evaporate. Isolate the desired  
 15 compound  
 (R<sub>f</sub> = 0.4 in 5% CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>) by column chromatography on silica  
 gel, and evaporate to a pale brown foam (0.54 g).


Convert a small portion to the HCl salt in the previously described  
 manner.

20 Mp: 240-250°C, with decomposition; Mass spectrum: MH<sup>+</sup> = 551.

-34-

Stir a mixture of free base (8) (0.5 g), EtOAc (5 mL) and conc. HCl (3 ml) at RT for 2h. Add excess aq. NaOH, extract with CH<sub>2</sub>Cl<sub>2</sub>, dry and evaporate to a foam (0.35 g). Dissolve 0.12 g of this in CH<sub>2</sub>Cl<sub>2</sub> (4 ml) and stir for 2h. at RT with 1N-NaOH (5 ml) and o-toluoyl chloride (0.05 mL). Separate and evaporate the organic phase, and isolate the product by preparative tlc on silica with 5% CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>. Convert a small portion to the hydrochloride in the usual way. Mp: 220-230°C, with decomposition; Mass spectrum: MH<sup>+</sup> = 569.

**Example 7: Synthesis of Compound No. 25**




10


Stir a solution of the product of example 6 (0.08 g) in DMF (1 ml) and add NaH (0.044 g), then after 10 min. add CH<sub>3</sub>I and stir for 30 min. Work up in EtOAc-H<sub>2</sub>O, dry, evaporate, purify by preparative tlc and convert to the HCl salt as in the foregoing preparation. Mp: 190-200°C, with decomposition; Mass spectrum: MH<sup>+</sup> = 583.

15

**Example 8: Synthesis of Compound No. 83 and 84**



-35-



300 ml of trifluoroacetic anhydride (TFAA) was added to 96 g of isonipecotic acid at 0°C. After the addition, the reaction mixture was

-36-

heated at reflux for 4h. Excess TFAA was removed under vacuo and the reaction mixture taken up in EtOAc and washed with water and concentrated to give 160 g of the amide. 50 g of this amide was treated with 300 ml  $\text{SOCl}_2$  and the reaction mixture heated at reflux overnight.

- 5 At the end of this time excess  $\text{SOCl}_2$  was removed under vacuo to give 54 g of the acid chloride.

- 11g  $\text{AlCl}_3$  was added slowly to a solution of 10 g of the above acid chloride in 40 ml of bromobenzene at ambient temperature and the reaction mixture heated at reflux for 4 h. It was then cooled and poured into a mixture of conc. HCl and ice and product extracted with EtOAc. The organic layer was separated and washed with water, half saturated  $\text{NaHCO}_3$  solution and concentrated to give 16.21 g of the ketone.

- 16.21 g of the ketone was dissolved in 200 ml toluene containing 25 ml ethylene glycol and 0.5 g p-toluenesulfonic acid. The reaction mixture was heated at reflux with azeotropic removal of water until no further water was collected. The reaction mixture was concentrated to give 17.4 g of the ketal.

- 17.4 g of the crude ketal was dissolved in 100 ml of  $\text{CH}_3\text{OH}$  and to this was added 25 ml of water and 12 g of  $\text{K}_2\text{CO}_3$  and the reaction mixture stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was separated and washed with water and brine and concentrated to give 12.55 g of (10).

- A mixture of 6.5 g of (10), 4.15 g of N-BOC-4-piperidone and 10 ml of  $\text{Ti}(\text{OiPr})_4$  was stirred at ambient temperature overnight. The reaction mixture was cooled to  $0^\circ\text{C}$  and 3.2 g of  $\text{NaCNBH}_3$  dissolved in 40 mL of  $\text{CH}_3\text{OH}$  was added. The reaction mixture was then allowed to warm to ambient temperature and stirred for 1 h., diluted with 100 ml of EtOAc and quenched with a mixture of 60 ml water/20 ml of conc.  $\text{NH}_4\text{OH}$ . The mixture was stirred for 1 h. and filtered through celite. The aqueous layer was extracted with 3x100 ml of EtOAc. The combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude was purified by flash chromatography using 100% EtOAc as eluent. To give 6.8 g, 66% of (11).



-37-

To a stirred solution of (11) in 25 mL of THF at -78°C was added 1.7 ml (1.6 M in Hex.) of <sup>n</sup>BuLi. The mixture was stirred at -78°C for 10 min., then 700 mg of piperonal dissolved in 5 ml of THF was added. The mixture was allowed to reach ambient temperature and stirred for 2 h.

- 5 The mixture was quenched with 30 ml of water and extracted with 3x50 ml of EtOAc. The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated. The crude was purified by silica gel flash chromatography using 10%Et<sub>3</sub>N-ether to give 475 mg, 41.5% of (12).


- 10 To a stirred solution of 475 mg of (12) and 1 ml of Et<sub>3</sub>SiH in 10 ml of CH<sub>2</sub>Cl<sub>2</sub> was added 3 ml of TFA. The mixture was stirred at ambient temperature for 2 h. The solvent was evaporated and the residue was taken up in 20 ml of CH<sub>2</sub>Cl<sub>2</sub>, washed with 20 ml of 10% NaOH. The aqueous layer was extracted with 3x30 ml of CH<sub>2</sub>Cl<sub>2</sub>, the combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated to give  
15 a mixture of (13) & (14) which was used without purification. (350 mg, 90%).

- 350 mg of the crude mixture of (13) and (14) and 1 ml of Et<sub>3</sub>N in 10 ml of CH<sub>2</sub>Cl<sub>2</sub> was treated with 0.5 ml (1.2 eq.) of o-toluoyl chloride for 3 h. The mixture was then concentrated and the residue purified by silica  
20 gel prep. TLC using 4% Et<sub>3</sub>N-Et<sub>2</sub>O to give compounds 83 and 84. The HCl salts of compounds 83 and 84 were prepared by dissolving the free base in EtOAc and adding HCl-ether solution. After evaporating the solvent, the salts were collected as powder.

- 25 Compound No. 83 (50 mg, 54%) (M+H)<sup>+</sup> Cal: 525.2753;  
Found: 525.2746

Compound No. 84 (190 mg, 50%) (M+H)<sup>+</sup> Cal: 569.3015;  
Found: 569.3022

-38-

Example 9: Synthesis of Compound No. 85

5 The piperazine (17) (1.1 g, 2.8 mmol), prepared as described in  
 example 1, is dissolved in  $\text{CH}_2\text{Cl}_2$  (10 ml) followed by addition of the N-  
 BOC piperidinone (0.85 g, 4.25 mmol) and  $\text{Ti}(\text{OiPr})_4$  (0.8 g, 2.8 mmol).  
 The mixture is stirred overnight at room temperature. A 1 molar toluene  
 solution of diethylaluminum cyanide (6.5 ml, 5.6 mmol) is added, via  
 10 syringe, and the mixture stirred for an additional 20h. The mixture is  
 then diluted with  $\text{EtOAc}$  and washed with 10%  $\text{Na}_2\text{CO}_3$ , water and brine.  
 The organic phase is concentrated in vacuo and the residue

-39-


chromatographed through a silica gel column (1:1 EtOAc/hexane). The product (18) is collected as a colorless oil (0.93 g, 55%).

The cyano intermediate (18) (0.13 g, 0.19 mmol) is dissolved in THF (4 ml) followed by the addition of  $\text{CH}_3\text{MgBr}$  (0.64 ml, 3.0 M, 1.9 mmol), via syringe. After stirring for 10 min the temperature is raised to 60°C where stirring is continued for 2.5h. Cooling to room temperature is followed by quenching with water (1 ml) and then saturated  $\text{NaHCO}_3$  (20 ml). After 10 min EtOAc (50 ml) is added, stirred vigorously, and separated. The organic layer is washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . Filtration and concentration with under vacuum gives a viscous oil which is purified on silica gel prep TLC plates (100% EtOAc). The intermediate (19) is collected in 60% yield.

The N-BOC piperidine (19) (0.27 g, 0.4 mmol) is dissolved in EtOAc (10 ml) and 1.5 ml of a 6N HCl solution added with vigorous stirring. After 1.5h saturated  $\text{NaHCO}_3$  is added until the pH was basic.  $\text{CH}_2\text{Cl}_2$  (25 ml) is added and the layers separated, and the aqueous phase extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer is dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to collect 0.22 g, 95%, of (20) as a thick oil which solidifies under vacuum.

The amine (20) (0.025 g, 0.044 mmol) is dissolved in  $\text{CH}_2\text{Cl}_2$  (1 ml) followed by the addition of  $\text{Et}_3\text{N}$  (11  $\mu\text{l}$ , 0.08 mmol) and the 2,6-difluoro-benzoyl chloride (0.014 g, 0.08 mmol). After stirring at room temperature overnight, saturated  $\text{NaHCO}_3$  and  $\text{CH}_2\text{Cl}_2$  are added. The layers are separated and the organic phase dried over  $\text{Na}_2\text{SO}_4$ . The solvent is removed and the residue dissolved in EtOAc. The solution was placed on a preparative TLC plate and eluted with EtOAc (100%). Compound 85 is collected as a clear oil (0.026 g), yield = 95%. This is converted to the dihydrochloride salt as follows: The free amine (26 mg) is dissolved in 1 ml EtOAc and stirred while HCl /  $\text{Et}_2\text{O}$  is added until the pH persists at 2. The precipitate is transferred to a centrifuge tube and spun in a centrifuge. The supernatant was removed and replaced with  $\text{Et}_2\text{O}$  (gentle stirring with a spatula), and the sample spun again. The process is repeated 3X. After removal of the solvent, the solid is transferred to a vial where it is dried under vacuum. 15 mg of a white solid was collected (m.p. = 242-245 dec).

-40-

**Example 10: Synthesis of Intermediate (3a) from example 3****Step 1**

- 5 TFAA (150 ml, 1.06 mol) in  $\text{CH}_2\text{Cl}_2$  (500 ml) was cooled to 0°C. A solution of (S)-α-methylbenzylamine (22) (100 g, 0.83 mol) in  $\text{CH}_2\text{Cl}_2$  (200 ml) was added slowly over 30 min. The cooling bath was removed and the mixture was stirred for ~2h, transferred to a separatory funnel,

-41-

washed with water, dried over anhydrous  $\text{MgSO}_4$ , filtered, and concentrated to give 189 g (~100%) of (23) as a white solid

**Step 2**

- Iodine (41 g, 162 mmol) was added to a solution of (23) (65.4 g, 301 mmol) in  $\text{CH}_2\text{Cl}_2$  (500 ml) at room temperature.
- [Bis(trifluoroacetoxy)iodo]-benzene (75 g, 174 mmol) was added. The mixture was stirred for 30 min, a mild exotherm occurred ( $30^\circ\text{C}$ ), then kept in the dark overnight. The mixture was added slowly to a stirred mixture of  $\text{NaHCO}_3$  (50 g), sodium bisulfite (10g) and water (600 ml), washing in with  $\text{CH}_2\text{Cl}_2$ . The mixture was stirred for 15 min, filtered and the filter cake was well washed with  $\text{CH}_2\text{Cl}_2$  until TLC (100%  $\text{CH}_2\text{Cl}_2$ ,  $R_f=0.75$ ) shows no product in the washings. (Note: the initial solid contains considerable product which is moderately soluble in  $\text{CH}_2\text{Cl}_2$ ). The filtrate was dried over anhydrous  $\text{MgSO}_4$ , filtered through a 1" pad of flash silica gel, washed with 2%  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ . The filtrate was evaporated to a slurry,  $\text{Et}_2\text{O}$  (~75 ml) and hexanes (~700 ml) were added and the mixture was stirred for 30 min. The resulting solid was collected, washed well with hexanes, air dried and then dried under vacuum to give 63.05 g (61%) of (24) as a white solid. Filtrates were evaporated to an oil and some crystals, added hexanes (500 ml), stirred at room temperature for 1h, collected and additional 4.6 g (4%) of (24)

**Step 3**

- (24) (10g, 29.15 mmol) was dissolved in  $\text{CH}_3\text{OH}$  (150 ml) and water (15 ml). 50%  $\text{NaOH}$  (30g) was added and the mixture was stirred at room temperature overnight (20 h). TLC (50%  $\text{EtOAc}/\text{hex.}$ ) indicated consumption of starting material. Most of  $\text{CH}_3\text{OH}$  was removed in vacuo, the residue was dissolved in water and  $\text{CH}_2\text{Cl}_2$ , transferred to a separatory funnel and extracted with  $\text{CH}_2\text{Cl}_2$ . The extracts were combined, dried over anhydrous  $\text{Na}_2\text{CO}_3$ , filtered and concentrated to give 6.7 g (93%) of (25) as a white solid.

**Step 4**

- Triflic anhydride (10.6 ml, 62.98 mmol) was added to a  $0^\circ\text{C}$  solution of (S) ethyl lactate (26) (7.0 ml, 61.75 mmol) in  $\text{CH}_2\text{Cl}_2$  (100 ml) at  $0^\circ\text{C}$ . 2,6-Lutidine (7.55 ml, 64.83 ml) was added and the mixture was stirred for 1h. The mixture was transferred to a separatory funnel,

-42-

washed with 0.5 M HCl and brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , concentrated almost to dryness, filtered through a plug of silica eluting with  $\text{CH}_2\text{Cl}_2$ , conc. to provide 10.8 g (70%) (27) as a reddish oil.

**Step 5**

5 (25) (6.7 g, 27.11 mmol) was added to a mixture of  $\text{CH}_2\text{Cl}_2$  (90 ml), water (90 ml) and  $\text{K}_2\text{CO}_3$  (5.5 g, 39.6 mmol). Ethyl (S) lactate triflate (27) (7.5 g, 29.82 mmol) was added and the mixture was stirred overnight (20 h). TLC (50% EtOAc/hex.) indicated consumption of starting material. Ammonia (conc.) (30 ml) was added, stirred 15 min,  
10 transferred to a separatory funnel, washed with water, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give 10.3 g (~110%) of (28).

**Step 6**

15  $\text{ClCH}_2\text{OCl}$  (12 ml, 150 mmol) was added to a solution of (28) (9.37g, 27.11) in dichloroethane (80 ml). The mixture was refluxed for 2h. TLC (20% EtOAc/hex.) indicated consumption of starting material. The solution was cooled to room temperature, diluted with  $\text{CH}_2\text{Cl}_2$ , water (200 ml) followed by  $\text{K}_2\text{CO}_3$  (~15g) added in small portions. The mixture was stirred for an additional 15 min., transferred to a separatory  
20 funnel, washed with water, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give 11.53g (~110%) of (29).

**Step 7**

(29) (27.11 mmol) was dissolved in dimethyl sulfoxide (DMSO) (120 ml). Concentrated aqueous ammonia (24 ml) and NaI (13 g) were  
25 added.  $\text{CH}_3\text{OH}$  (10 ml) was used in the NaI. The mixture was stirred over the weekend (60h, overnight is sufficient). TLC (50% EtOAc/hex.) indicated consumption of starting material. Water (500 ml) was added, the mixture was stirred for 30 min and the resulting precipitate was collected via vacuum filtration. The precipitate was well washed with  
30 water and dried under vacuum to afford 8.2g (84%) of (30).

**Step 8**

(30) (8.2 g, 22.9 mmol) was added to a mixture of  $\text{NaBH}_4$  (8.66 g, 228.9 mmol) in dimethoxyethane (250 ml). Boron trifluoride etherate (16.9 ml, 137.3 mmol) was added and the resulting mixture was refluxed  
35 under nitrogen for 3h. TLC (5%  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) indicated consumption

-43-


of starting material. The mixture was cooled to 0°C, CH<sub>3</sub>OH (60 ml) was added slowly and the resulting mixture was stirred for 20 min.

Concentrated HCl (10 ml) was added slowly and the resulting mixture was refluxed for 1h. TLC (5% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) indicated formation of a more polar product consistent with an amine. The mixture was cooled to room temperature and concentrated almost to dryness. The residue was partitioned between 2N NaOH and CH<sub>2</sub>Cl<sub>2</sub>, transferred to a separatory funnel and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extracts were combined, washed with water, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated to provide 6.5 g (87%) of (31) as an oil.

**Step 9:**

Intermediate (31) was converted to 3a using using procedures described in example 8.

**Example 11: Synthesis of Intermediate (3b) from example 3**




A solution of TFAA (0.4 mol) in 300 ml cold CH<sub>2</sub>Cl<sub>2</sub> is treated with 0.3 mol (S)-α-methylbenzylamine in 100 ml CH<sub>2</sub>Cl<sub>2</sub>. The resulting mixture is stirred with ice cooling for 30 minutes and at room temperature for 1.5 hours. The solution is again cooled in an ice bath and 80 ml CH<sub>3</sub>SO<sub>3</sub>H is added followed by 45 grams dibromodimethylhydantoin. Stirring is continued until the mixture is homogeneous, then the mixture is allowed to stand overnight in the dark. A solution of 20 g NaHSO<sub>3</sub> in 300 ml ice water is added. The organic layer is separated and washed twice with water and dried over MgSO<sub>4</sub>. The solution is filtered through a 4" pad of flash-grade silica gel, eluting with 400 ml CH<sub>2</sub>Cl<sub>2</sub>, and

-44-

evalaporated to a solid. The solid is suspended in 200 ml Et<sub>2</sub>O and 1l hexane is added. After stirring 30 minutes, the solids are collected, washed with hexane, and dried to give 35.7 grams of intermediate (24b). This compound is converted to (3b) using the same sequence described in scheme 9.

### Example 12: Synthesis of Compound No. 53



To a stirred solution of (32) prepared via the procedure described for intermediate (7) of example 14, but using N-ethoxycarbonyl-4-piperidinone instead of N-tBOC-4-piperidinone (1.80 g, 3.41 mmol) in toluene (34 ml) was added dry ethylene glycol (1.33 ml, 23.8 mmol), triethylorthoformate (1.70 ml, 10.2 mmol) and *p*-toluenesulfonic acid monohydrate (0.97 g, 5.12 mmol). The reaction was heated overnight at 55°C under nitrogen. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed sequentially with 1N NaOH and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and




-45-

concentrated in vacuo. The resulting product (33) (2.40 g) was used without further purification.

The crude ketal (33) (0.83 g, ~1.45 mmol) was dissolved in ethylene glycol (44 ml) and crushed KOH (2.90 g, 51.6 mmol) was added to the rigorously stirred solution. The reaction was heated for 24 h at 100°C. After cooling to room temperature, the mixture was diluted with EtOAc followed by the addition of 1N NaOH. The aqueous portion was extracted with EtOAc and the combined organics washed several times with 1N NaOH. The crude solution was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield 0.31 g of (34) which was used without further purification.

To a solution of (34) (31.4 mg, 62.8 µmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.60 ml) was added *n*-propylsulfonyl chloride (20 µL, 178 µmol) and Et<sub>3</sub>N (30 µL, 215 µmol). The reaction was stirred for 2 h at room temperature under nitrogen. The mixture was concentrated under reduced pressure and purified by PTLC (10/90 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>), yielding (35) (12.3 mg, 32% over 3 steps). LRMS (FAB): (M+H)=606

### Example 13: Synthesis of Compound No. 22




To a solution of (36) prepared via the procedure described for intermediate (7) of example 14, substituting 4-mercaptoanisole for 4-mercaptomethylenedioxybenzene and N-ethoxycarbonyl-4-piperidinone for N-tBOC-4-piperidinone (29.5 mg, 58.9 µmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.24 ml) was added ethanedithiol (7.4 µL, 88 µmol) and BF<sub>3</sub>·OEt<sub>2</sub> (11 µL, 88 µmol). The reaction was stirred overnight at room temperature under nitrogen, then diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with 10% NaOH.

-46-

The crude mixture was dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated under reduced pressure. Purification by PTLC (10/90  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) yielded **37** (17 mg, 50% yield). LRMS (FAB):  $(\text{M}+\text{H})=591$ .

**Example 14: Synthesis of Compound 43**



10 To a stirred solution of 41 ml of anhydrous DMF was added 43 ml of  $\text{SO}_2\text{Cl}_2$  at  $0-10^\circ\text{C}$  followed by slow addition of 61 grams of methylenedioxybenzene. After the addition was completed, the mixture was heated at  $80^\circ\text{C}$  for 10 min., and then 5-10 min. at  $110^\circ\text{C}$ . The reaction mixture turned to dark brown. The mixture was cooled to  $40^\circ\text{C}$  and was poured into a mixture composed of 200 ml  $\text{H}_2\text{O}$ /200g crushed ice/300 ml  $\text{CHCl}_3$ . Two layers were separated, the organic layer was

-47-

dried over  $\text{MgSO}_4$ , filtered, and concentrated to give a crude product (1) which was triturated with 100 ml of hexanes. (60.6 g, 53%)

To a stirred suspension of 16 g LAH in 500 ml of THF at  $0^\circ\text{C}$ , was added dropwise 60.6 g of (1) dissolved in 100 ml of THF. The  
5 temperature was allowed to reach RT and the mixture was stirred at this temperature for 2 h.. The reaction mixture was then quenched carefully with 1N HCl at  $0^\circ\text{C}$  to pH=1, diluted with 1 L EtOAc. The organic phase was washed with 500 ml of water, dried over  $\text{MgSO}_4$ , filtered and concentrated to give a crude which was used directly in the next step  
10 without purification. (29 g, 68%)

28 g of the crude thiol (2) dissolved in 20 ml of DMF was added to a stirred suspension of 8 g of NaH(60%) in 80 ml of DMF. The mixture was stirred at RT for 1 h., ketone (3) dissolved in 150 ml of DMF was added at once. The mixture was heated at  $70^\circ\text{C}$  over night. The mixture  
15 was diluted with 400 ml of  $\text{CH}_2\text{Cl}_2$  and then quenched with 300 ml of water. The organic phase was washed with 3x200 ml of water, dried with  $\text{MgSO}_4$ , filtered and concentrated to give crude (4) which was purified by flash chromatography using EtOAc/Hex./ $\text{CH}_2\text{Cl}_2$ =1/4.5/4.5 as eluents. (35 g, 53%)

20 To a stirred solution of 6 g of (4) in 100 ml of  $\text{CH}_2\text{Cl}_2$  was added 10 ml of TFA. The mixture was stirred at RT for 1 h. The solvent was eliminated under reduced pressure and the residue was taken up in 100 ml of  $\text{CH}_2\text{Cl}_2$  and extracted with 3x60 ml of  $\text{CH}_2\text{Cl}_2$ . The combined organic phase was chloride, quenched with 50 ml of 10% NaOH. The  
25 aqueous phase was dried over  $\text{MgSO}_4$ , filtered and concentrated to give crude (5), used without purification. (4.6 g, 100%)

A mixture of 3.4 g of (5) 2.4 g of N-BOC-4-piperidone and 5.9 ml of  $\text{Ti}(\text{OiPr})_4$  was stirred at RT over night. 1.6 g of  $\text{NaCNBH}_3$  dissolved in 40 ml of  $\text{CH}_3\text{OH}$  was added at  $0^\circ\text{C}$ . The mixture was stirred at RT for 1 h.  
30 diluted with 100 ml of EtOAc, quenched with a mixture of 60 ml water/20 ml of  $\text{NH}_4\text{OH}$ . The mixture was stirred at RT for 1 h. and filtered though celite. The aqueous phase was extracted with 3x100 ml of EtOAc. The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude (6) was purified by flash chromatography using  
35 100% EtOAc as eluent. (3.8 g, 73%)


-48-

To a solution of 1.32 g of (6) in 15 ml of HOAc was added 1.41 g of  $\text{NaBO}_3 \cdot 4\text{H}_2\text{O}$ , and the mixture was stirred at RT for 3 days. The solvent was evaporated and the residue was quenched with 20 ml of 10% NaOH, then extracted with 3x50 ml of EtOAc. The combined organic extract was dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude (7) was purified by silica gel prep TLC using 20% EtOH-EtOAc. (620 mg, 44%)

620 mg of (7) in 15 ml of  $\text{CH}_2\text{Cl}_2$  and 5 ml of TFA was stirred at RT for 3 h. The solvent was evaporated and residue was treated with 10% NaOH, then extracted with EtOAc. The organic extract was dried over  $\text{MgSO}_4$ , filtered and concentrated to give crude product (510 mg, 100%). 0.052 mmol of the resultant crude free amine was reacted with 1.2 eq. of o-toluoyl chloride and 1.3 eq. of  $\text{Et}_3\text{N}$  in 3 ml of  $\text{CH}_2\text{Cl}_2$  for 3 h. The mixture was concentrated and the residue was purified by silica gel preparative TLC using 20% EtOH-EtOAc. (18 mg, 50%)  $(\text{M}+\text{H})^+$  Calcd: 575.2216; Found: 575.2223.

The HCl salts were prepared by dissolving 43 in EtOAc and adding HCl-ether solution. After evaporating the solvent, the salts were collected as powder.

**Example 15: Preparation of Compound No. 82**




-49-


Stir a solution of compound (4) from Example 3 (0.5 g) and  $\text{NaBH}_4$  (0.2 g) in ethanol (3 ml) at RT for 3 h. Add water, extract with  $\text{CH}_2\text{Cl}_2$ , dry over  $\text{MgSO}_4$  and evaporate. To the residue in  $\text{CH}_2\text{Cl}_2$  (5 ml) add triethylsilane (3 ml) and TFA (3 ml), reflux for 3 h., evaporate, partition with  $\text{CH}_2\text{Cl}_2$  and 1N aq. NaOH, dry and evaporate to obtain (11) as a thick oil, used as such in subsequent acylation experiments.

Stir a mixture of (11) (0.03 g),  $\text{CH}_2\text{Cl}_2$  (2 ml) and 1N aq. NaOH (2 ml) and add 2,3-dimethylbenzoyl chloride (0.07 g). Stir at RT for 18 h., extract with  $\text{CH}_2\text{Cl}_2$ , dry and evaporate. Dissolve the residue in 1:1  $\text{CH}_2\text{Cl}_2$ :acetone and filter through a small plug of silica gel, washing with the same solvent mixture. Evaporate, dissolve the residue in a small volume of  $\text{CH}_2\text{Cl}_2$  and precipitate the dihydrochloride by adding to excess HCl in  $\text{Et}_2\text{O}$  as described in earlier preparations. Wash with  $\text{Et}_2\text{O}$  and dry to obtain the product compound No. 82 as an off-white powder (0.036 g). Mp: 207-212°C, with decomposition; MS:  $\text{MH}^+ = 554$ .

#### Example 16: Preparation of Compound No. 94



-50-



To a stirred solution of 899 mg (1.72 mmol) of (4), prepared as described in Example 14, in 15 ml of anhydrous  $\text{CH}_3\text{OH}$  is added by portion 130 mg of  $\text{NaBH}_4$ . The mixture is stirred at room temperature for 30 min. and then quenched with 15 ml of 10%  $\text{NaOH}$  solution. The aqueous layer is extracted with 4x20 ml of  $\text{EtOAc}$ . The combined organic layers are dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give 859 mg (95%) of (8).

To a stirred solution of 850 mg of crude (8) in 5 ml of  $\text{CH}_2\text{Cl}_2$  is added 563 mg (3 eq.) of triethylsilane, followed by 2 ml of TFA. The mixture is stirred at room temperature for 3 hrs. and then is concentrated. The residue is taken up in 15 ml of  $\text{CH}_2\text{Cl}_2$  and 15 ml of 10%  $\text{NaOH}$ . The aqueous layer is extracted with 3x15 ml of  $\text{CH}_2\text{Cl}_2$ . The combined organic layers are dried over  $\text{MgSO}_4$ , filtered and concentrated to give 600 mg (92%) of crude (9).






To a stirred solution of 600 mg of (9) and 455 mg of  $\text{Et}_3\text{N}$  in 5 ml of  $\text{CH}_2\text{Cl}_2$  is added 464 mg of o-toluoyl chloride. The mixture is stirred at room temperature for 2 hrs. The mixture is quenched with 20 ml of water, extracted with 3x20 ml of  $\text{EtOAc}$ . The combined organic layers are dried over  $\text{MgSO}_4$ , filtered and concentrated to give a crude. Purification on silica gel prep TLC affords 555 mg (70%) of (10).

To a stirred solution of 43 mg of (10) and 0.5 ml of  $\text{CH}_3\text{SO}_3\text{H}$  (0.5 M  $\text{CH}_2\text{Cl}_2$ ) in 5 ml of  $\text{CH}_2\text{Cl}_2$  is added at  $0^\circ\text{C}$ , 58 mg (57-86%) of mCPBA. The mixture is stirred at  $0^\circ\text{C}$  for 1 h. and is quenched with 15 ml of 10%  $\text{NaHCO}_3$ . The aqueous layer was extracted with 3x15 ml of  $\text{EtOAc}$ . The combined organic layers are dried over  $\text{MgSO}_4$ , filtered and concentrated to give a crude product. Purification on silica gel prep. TLC affords 36 mg (80%) of free base of compound number 94. The HCl salt is prepared by dissolving the free base in  $\text{EtOAc}$  and adding HCl-ether solution. After evaporating the solvent, the salts are collected as powder.  $(\text{M}+\text{H})^+$  Cal: 561.2423; Found: 561.2416.







-51-

Using appropriate starting materials in the above procedures, with modifications well known in the art, the compounds exemplified in the following table are prepared:

TABLE OF COMPOUNDS







NO.	STRUCTURE	PHYSICAL DATA
<u>1</u>		-
<u>2</u>		-
<u>3</u>		-
<u>4</u>		mp=215-225°C (dec) MS MH <sup>+</sup> =574
<u>5</u>		-

-52-







<u>6</u>		-
<u>7</u>		-
<u>8</u>		-
<u>9</u>		-
<u>10</u>		-
<u>11</u>		-









-53-

<b>12</b>		-
<b>13</b>		-
<b>14</b>		-
<b>15</b>		-
<b>16</b>		-
<b>17</b>		mp=182-190 oC (dec) MS MH+=588

-54-







<b>18</b>		-
<b>19</b>		-
<b>20</b>		-
<b>21</b>		-
<b>22</b>		LRMS (FAB) Calc. for C <sub>29</sub> H <sub>38</sub> O <sub>5</sub> N <sub>2</sub> S <sub>3</sub> (M+H): 591 Found: 591
<b>23</b>		-

-55-







<u>24</u>		mp=220-230°C (dec) MS MH+=569
<u>25</u>		mp=190-200°C (dec) MS MH+ = 583
<u>26</u>		-
<u>27</u>		HRMS Calcd: 573.2278 Found: 573.2271
<u>28</u>		mp = 200-210°C (dec) MS MH+ = 554
<u>29</u>		-

<b>30</b>		mp = 85-95°C (dec) MS MH+ = 556
<b>31</b>		mp= 210-215°C (dec) MS MH+=540
<b>32</b>		mp=235-237°C (dec)
<b>33</b>		-
<b>34</b>		mp=185-200°C (dec) MH+ 608/610
<b>35</b>		mp=178-190°C (dec) MS MH+ = 652/654







-57-

<u>36</u>		mp=180-190°C (dec) MH+= 604
<u>37</u>		mp=180-195°C (dec) MH+=593
<u>38</u>		-
<u>39</u>		mp= 190-205°C (dec) MS MH+=578
<u>40</u>		-
<u>41</u>		mp = 180-290°C (dec) MS MH+=594







-58-

<b>42</b>		-
<b>43</b>		HRMS Calcd 575.2216 Found 575.2223
<b>44</b>		HRMS Calcd 589.2372 Found 589.2379
<b>45</b>		mp=255-257°C (dec)
<b>46</b>		FAB MH+=694
<b>47</b>		-

-59-






<b>48</b>		-
<b>49</b>		FAB MH+=632
<b>50</b>		mp=195-200°C (dec)
<b>51</b>		-
<b>52</b>		-
<b>53</b>		LRMS (FAB) Calc. for C <sub>29</sub> H <sub>38</sub> O <sub>8</sub> N <sub>2</sub> S <sub>2</sub> (M+H): 606 Found: 606

-60-







<b>54</b>		FAB (NBA-G/TG-DMSO): 590 (M+), 574, 406, 389
<b>55</b>		HRMS calcd 594.2672 found 594.2668
<b>56</b>		HRMS calcd 642.2686 found 642.2685
<b>57</b>		HRMS calcd 606.3002 found 606.2987
<b>58</b>		mp=218-220°C (dec)
<b>59</b>		mp=203-205°C (dec)









-61-

<u>60</u>		-
<u>61</u>		-
<u>62</u>		LRMS (FAB) Calc. for $C_{31}H_{40}O_8N_2S$ (M+H): 601 Found: 601
<u>63</u>		FAB MH+=682
<u>64</u>		HRMS calcd 636.3107 found 636.3119







-62-

<b>65</b>		-
<b>66</b>		mp=157-160°C (dec)
<b>67</b>		FAB MH+=632
<b>68</b>		mp=172-175°C (dec)
<b>69</b>		
<b>70</b>		







-63-

<b>71</b>		
<b>72</b>		
<b>73</b>		
<b>74</b>		
<b>75</b>		
<b>76</b>		







-64-

<u>77</u>		
<u>78</u>		
<u>79</u>		HRMS calcd 622.2951 found 622.2932
		HRMS calcd 606.3002 found 606.2999
<u>81</u>		HRMS calcd 634.2952 found 634.2962
<u>82</u>		mp: 207-212 °C (dec) Mass spectrum: MH+ = 554







-65-

<b>83</b>		-
<b>84</b>		HRMS Calcd 579.3015 Found 569.3022
<b>85</b>		mp=242-245 (dec)
<b>86</b>		mp=206-209 (dec)
<b>87</b>		mp=237-239 (dec)
<b>88</b>		FAB (SIMS): 630 (M+), 391, 329, 307, 289







-66-

<b>89</b>		FAB (SIMS): 568 (M+), 391, 307, 232
<b>90</b>		FAB (NBA-G/TG- DMSO): 554 (M+), 538, 389, 289
<b>91</b>		FAB (NBA-G/TG- DMSO): 610 (M+), 594, 490, 426, 322
<b>92</b>		HRMS calcd 572.3488 found 572.3491
<b>93</b>		HRMS calcd 620.2488 found 620.2478
<b>94</b>		HRMS Calcd 561.2323 Found 561.2423

-67-







<b>95</b>		<p>HRMS calcd 700.2056 found 700.2044</p>
<b>96</b>		<p>LRMS (FAB) Calc. for C<sub>33</sub>H<sub>35</sub>O<sub>7</sub>N<sub>2</sub>SCl (M+H): 639 Found: 639</p>
<b>97</b>		<p>FAB (NBA-G/TG-DMSO): 616 (M+), 432, 389, 328</p>
<b>98</b>		-
<b>99</b>		-
<b>100</b>		<p>mp: 210-220°C (dec) Mass spectrum: - MH<sup>+</sup> = 546.</p>

-68-




<b>101</b>		-
<b>102</b>		HRMS calcd 650.2593 found 650.2588
<b>103</b>		HRMS calcd 606.3099 found 606.3084
<b>104</b>		HRMS calcd 592.3209 found 592.3215
<b>105</b>		HRMS calcd 592.3209 found 592.3215
<b>106</b>		HRMS calcd 606.3099 found 606.3090




-69-


<u>107</u>		HRMS calcd 650.2593 found 650.2579
<u>108</u>		-
<u>109</u>		-
<u>110</u>		LRMS(FAB): M+H = 732
<u>111</u>		LRMS(FAB): M+H = 724
<u>112</u>		mp: 190-195 °C (dec) Mass spectrum: MH+ = 546.

-70-

<b>113</b>		$MH^+ = 618, 620$
<b>114</b>		mp=191°C (dec) $MH^+ = 611$
<b>115</b>		$MH^+ = 540$

**Example 17**

Step 1:




- 5 To a mixture of piperidone monohydrate hydrochloride (27.5 g, 0.18 mol),  $K_2CO_3$  (50g), 350 ml  $CH_2Cl_2$ , 250 ml water, cooled in an ice water bath, was added dropwise (over 1h) a solution of o-toluoyl

-71-

chloride (26 g, 0.17 mol) in 25 ml  $\text{CH}_2\text{Cl}_2$ . The mixture was warmed to room temperature and stirred overnight. The organic phase was separated, washed with water, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated on a rotary evaporator. The crude product was purified on a silica column using hexane: EtOAc (4:1). Yield = 93%

Step 2:




To a solution of 2-(R)-methylpiperazine (6.1 g, 61 mmol) in  $\text{CH}_2\text{Cl}_2$  (250 ml) was added acetic acid (3.6 ml), N-o-toluoypiperidone (13.3 g, 61 mmol), and  $\text{NaBH}(\text{OAc})_3$  (15.5 g, 73 mmol). The mixture was stirred overnight followed by dilution with  $\text{CH}_2\text{Cl}_2$ , washing with 10%  $\text{Na}_2\text{CO}_3$  and water. Concentration of the organic phase was followed by purification on a silica column ( $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH} = 20:1$ , sat'd with aq.  $\text{NH}_3$ ). Yield = 50%

Step 3:

To  $\text{NaBH}(\text{OAc})_3$  (0.042 g, 0.2 mmol) was added a solution of the product of Step 2 (0.03 g, 0.1 mmol), acetic acid (0.006 ml), and 4-phenoxy-benzaldehyde (0.1 mmol). The mixture was stirred overnight followed by dilution with  $\text{CH}_2\text{Cl}_2$  (10 ml), washing with 10%  $\text{Na}_2\text{CO}_3$ , and drying over  $\text{Na}_2\text{SO}_4$ . Concentration of the organic phase was followed by purification on a silica column (EtOAc). The hydrochloride was prepared by dissolving the product in a minimum amount of EtOAc, followed by addition of anhydrous HCl. The precipitate was collected on a centrifuge and washed with  $\text{Et}_2\text{O}$  (3x), followed by drying under vacuum. Yield = 66%. M.p. 217-219°C (dec).

#### Example 18




**117**


To  $\text{NaBH}(\text{OAc})_3$  (0.026 g, 0.12 mmol) was added  $\text{CH}_2\text{Cl}_2$  (1 ml), a solution of cyclohexanecarboxaldehyde (0.125 ml, 1M), acetic acid

-72-

(0.004 ml), and intermediate (20) of Example 9 (0.03 g, 0.06 mmol). The mixture was stirred overnight followed by dilution with  $\text{CH}_2\text{Cl}_2$  (10 ml), washing with 10%  $\text{Na}_2\text{CO}_3$  and drying over  $\text{Na}_2\text{SO}_4$ . Concentration of the organic phase was followed by purification on a silica column (EtOAc). The hydrochloride was prepared as described in Example 17. Yield=67%. M.p. 208-209°C (dec).

**Example 19**

160 mg. of compound **28** were taken up in 4 ml ethanol with 45 mg of  $\text{H}_2\text{NOH}\cdot\text{HCl}$ . 2 ml of pyridine were added and the mixture was heated to reflux for 5 hours. The solvent was removed *in vacuo*, and the reaction was diluted with  $\text{CH}_2\text{Cl}_2$  and extracted with aqueous  $\text{NaHCO}_3$ . The organic layers were dried over  $\text{MgSO}_4$ , evaporated, and the residue purified by column chromatography in a gradient of  $\text{CH}_3\text{OH}/\text{EtOAc}$ . This gave 130 mg of product as a mixture of *syn* and *anti* isomers. FAB ( $\text{MH}^+$ ) = 541.


**Examples 20A and 20B****20A:**

1.1 g of the product of Example 19 were taken up in 15 ml  $\text{CH}_2\text{Cl}_2$  with 2 eq. of  $\text{Et}_3\text{N}$ , and cooled to  $-78^\circ\text{C}$  under  $\text{N}_2$ . 440 mg (1.1 eq.) of chlorodiphenylphosphine were taken up in 2 ml of  $\text{CH}_2\text{Cl}_2$  and added slowly to the reaction at this temperature. The reaction was stirred for 2 h in the cold and allowed to come to room temperature overnight. The solvent was removed *in vacuo*, and the crude material was taken up in 20 ml THF. 100 mg (excess)  $\text{NaBH}_4$  was added and the mixture was stirred for 16 h at room temperature. The solvent was removed *in vacuo*, and the crude product was dissolved in a mixture of 10 ml of 6N HCl and


-73-

30 ml of CH<sub>3</sub>OH. This mixture was stirred overnight at room temperature. The CH<sub>3</sub>OH was evaporated from the mixture, and the reaction was taken to pH 9 with the addition of aqueous NaOH. This solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the organic layers washed with water, dried over MgSO<sub>4</sub> and evaporated to a crude oily product. This was purified by column chromatography in a gradient of CH<sub>3</sub>OH/EtOAc to give 356 mg of amine product. MS (FAB): 527.5 (M+1), 510.5

20B:

**120**

- 25 mg of the product of Step 20A were taken up in 2 ml CH<sub>2</sub>Cl<sub>2</sub> with 90 mg of sodium triacetoxyborohydride. 9 microliters of acetaldehyde were added, and the mixture was stirred at room temperature for 1.5 hours. The reaction was diluted with additional solvent and extracted with aqueous NaHCO<sub>3</sub>. The organic layers were dried over MgSO<sub>4</sub>, evaporated, and purified by preparative thin-layer chromatography in 5% CH<sub>3</sub>OH/EtOAc. This gave 12 mg of the desired product as an oil. FAB (M+) = 583.


**Example 21****121**

- 44 mg. of **28** were taken up in 20 ml of dry toluene with 2 ml of ethylene glycol and excess p-toluenesulfonic acid. The reaction was heated to reflux with a Dean-Stark water trap attached for 24 hours. NMR analysis showed partial reaction. Additional glycol and acid were added, and the reaction was heated an additional 24 hours. The mixture was extracted with aqueous NaHCO<sub>3</sub>, evaporated, and purified by column chromatography in a gradient of CH<sub>3</sub>OH/EtOAc to give 15 mg of




-74-

product, which still contains small amounts of the starting ketone.

MS (electrospray): 598 (M+1), 302.

**Example 22****122**

- 5            24 mg of compound **30** were taken up in 1 ml of  $\text{CF}_3\text{CO}_2\text{H}$ . An excess of 2-oxazolidinone was added and the mixture was stirred for 72 h at room temperature. Excess saturated  $\text{NaHCO}_3$  solution was added, and the mixture was extracted with EtOAc. This was dried over  $\text{Na}_2\text{SO}_4$ , evaporated, and the residue purified by column chromatography in 5%  $\text{CH}_3\text{OH}/\text{EtOAc}$  to give approx. 15 mg of product as an oil.
- 10           MS (electrospray): 625.2 (M+1), 302.


**Example 23****123****124**Step 1:

15


- 11 g of piperidinone hydrochloride were stirred in 100 ml  $\text{CH}_2\text{Cl}_2$ . 100 ml of 20% aqueous  $\text{K}_2\text{CO}_3$  solution were added with vigorous stirring, followed by 8.8 ml of ortho-toluyyl chloride. The mixture was stirred overnight at room temperature. The organic layer was separated and the aqueous layer was washed with additional solvent. The combined  $\text{CH}_2\text{Cl}_2$  layers were dried over  $\text{Na}_2\text{SO}_4$ , evaporated, and purified by column chromatography in a gradient of hexane/EtOAc to give the product as an oil that solidified on standing.
- 20

Step 2:


-75-



- 5.55 g of the product of Step 1 were combined with 4.75 g (1 eq.) of N-(t-butoxycarbonyl)piperazine in 100 ml of  $\text{CH}_2\text{Cl}_2$ . Sodium triacetoxyborohydride (5.4 g) was added and the reaction was stirred at room temperature for 72 hours. An additional gram of sodium triacetoxyborohydride was added, along with 2 ml of acetic acid, and the mixture was stirred an additional 24 hours. 25 ml of  $\text{CH}_3\text{OH}$  was added and the mixture was evaporated. The crude material was dissolved in  $\text{CH}_2\text{Cl}_2$  and extracted with water. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , evaporated, and the residue purified by column chromatography in a gradient of EtOAc/hexane to give 6.7 grams of product.

Step 3:

- 6.7 g of the product of Step 2 were taken up in 100 ml  $\text{CH}_2\text{Cl}_2$ , and 20 ml of  $\text{CF}_3\text{CO}_2\text{H}$  were added slowly at room temperature. The reaction was stirred for 24 hours. 20 ml of water were added and the reaction was brought to pH 10 by the slow addition of solid NaOH. The organic layers were separated and the aqueous layers washed with additional  $\text{CH}_2\text{Cl}_2$ . The organic layers were dried over  $\text{Na}_2\text{SO}_4$  and evaporated to give 5.1 g of product as a solid, which was used directly.

Step 4:

- 3.9 g of the product of Step 3 were taken up in 200 ml THF with 7.15 g (3 eq.) of 1,4-(bis)-chloromethylbenzene and 2 ml (1 eq.) of  $\text{Et}_3\text{N}$ . The mixture was heated to reflux for 7 hours and cooled to room

-76-



temperature. 300 ml of Et<sub>2</sub>O were added and the precipitated Et<sub>3</sub>N•HCl was removed by filtration. 34 ml of 1M HCl in Et<sub>2</sub>O (2.1 eq.) was added in portions, with stirring, and the precipitate was filtered off and dried in vacuo to give 5.6 g of the dihydrochloride salt of the product, slightly contaminated with Et<sub>3</sub>N•HCl.

**Compound 123:**

65 mg of the product of Step 4 (as a mixture containing 60% Et<sub>3</sub>N•HCl) were taken up in 2 ml dry DMF with excess K<sub>2</sub>CO<sub>3</sub>. The mixture was stirred for 1 h at room temperature until the solid starting material had dissolved. In a separate flask, excess succinimide was reacted with NaH in 1 ml dry DMF for 30 min. at room temperature. The DMF solution of the free amine was removed from the K<sub>2</sub>CO<sub>3</sub> and added to this anion, and the mixture was stirred for 16 h at room temperature. Water was added, and the mixture was extracted twice with EtOAc. The organic layers were washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. This gave the product in sufficient purity to form its HCl salt directly. MS (electrospray): 489.1 (M+1), 276.

**Compound 124:**


In a manner similar to that described for compound 123, but using pyrrolidinone in place of succinimide and potassium hexamethyldisilazide (KHMDS) (in toluene) in place of NaH, compound **124** was prepared in sufficient purity to form its HCl salt directly. MS (electrospray): 475 (M+1), 408.

**Example 24:****Step 1:**



-77-

- 0.8 ml (6.35 mmol) of N-methyl-o-toluidine, 1.0 g (5.29 mmol) of 4-chloromethyl(benzoyl chloride) and 1.4 ml (7.93 mmol) of (iPr)<sub>2</sub>NEt were taken up in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) at 0°C. The solution was warmed to r.t., stirred at r.t. for 17 h, then was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with 1M HCl(aq.). The aqueous layer was washed with CH<sub>2</sub>Cl<sub>2</sub>. The combined CH<sub>2</sub>Cl<sub>2</sub> layers were dried over NaSO<sub>4</sub>, filtered and concentrated to give the desired benzyl chloride as a light brown oil (1.55 g, quant.).

Step 2:

- The product of Step 1 (1.45 g, 5.29 mmol), piperazine-piperidine Boc compound (1.42g, 5.29 mmol), and (iPr)<sub>2</sub>NEt (1.84 ml, 10.6 mmol) were taken up in CH<sub>3</sub>CN (25 ml). The solution was heated at reflux for 2.75 h. The solution was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 1 N NaOH (aq.). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined CH<sub>2</sub>Cl<sub>2</sub> layers were dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and concentration gave a tan foam (2.7 g). Purification via flash chromatography(15/1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH, SiO<sub>2</sub>) gave 2.26 g (84 %) of a colorless foam.

Step 3:


- The BOC-protected product of Step 2 (2.26 g, 4.47 mmol) was taken up in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) and cooled to 0°C. TFA (5 ml) was added, and the solution was warmed to r.t. The solution was stirred at r.t. for 20 h. The solution was concentrated and the residue triturated with CH<sub>2</sub>Cl<sub>2</sub>/Et<sub>2</sub>O to give a white solid. The solid was washed with Et<sub>2</sub>O and dried, which gave 4.1 g (99 %) of the TFA salt of the amine.

Step 4:

Using a procedure similar to that described in Example 1, treat the product of Step 3 to obtain **125**. M.p. = 235 - 240°C. HRMS (FAB) calc'd for C<sub>33</sub>H<sub>41</sub>N<sub>4</sub>O<sub>2</sub> (MH): 525.3230. Found: 525.3239.

-78-

## Example 25



- 5 **Step 1:** To a stirred solution of 10.0 g (22.6 mmol) of (61) in 50 ml of anhydrous  $\text{CH}_3\text{OH}$  was added  $\text{NaBH}_4$  in portions at  $0^\circ\text{C}$ . The mixture was stirred at rt for 30 min and then quenched with water. The aqueous layer was extracted with (4x30 ml)  $\text{EtOAc}$ . The combined layers were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give 10.0 g (99%) of an alcohol which was used directly. To a stirred solution of the alcohol 10.0 g (22.5 mmol) in  $\text{CH}_2\text{Cl}_2$  (100 ml) was added 16 ml of  $\text{Et}_3\text{SiH}$ , followed by 40 ml of TFA. The mixture was stirred at rt for 3 h and then quenched with 1N  $\text{NaOH}$ . The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (100 ml). The combined organic layers were dried over  $\text{Na}_2\text{SO}_4$  and concentrated to yield 6.0 g (82%) of (62) as a colorless oil.
- 10
- 15

-79-

**Step 2:** To a stirred solution of (62) (5.0 g, 15.29 mmol) in Et<sub>2</sub>O (200 ml) and 40 ml of 10% NaOH was added BOC<sub>2</sub>O. The mixture was then stirred at rt for 24 h. The layers were separated and the organic layer washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and chromatographed to yield 5.7 g (87%) of (63) as colorless oil.

**Step 3:** To a stirred solution of (63) (5.7 g, 13.34 mmol) in acetic acid was added NaBO<sub>3</sub>•4H<sub>2</sub>O and stirred for 24 h at rt. The reaction mixture was then quenched with 25% NaOH. The organic layer was then extracted with EtOAc (3x30 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and chromatographed to yield 2.8 g (46%) of (64).

**Step 4:** To a stirred solution of (64) (1.5 g, 3.26 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (35 ml) was added TFA (2 ml). The mixture was stirred at rt for 1 h and concentrated. The residue was then diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with 10% NaOH. The extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to yield 0.983 g (84%) of (65).

**Step 5:** To a stirred solution of (65) (0.983 g, 2.73 mmol) and t-butoxy-carbonyl piperidine (0.571 g, 2.87 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added Ti(iOPr)<sub>4</sub> (1 ml). The reaction mixture was stirred at rt for 24 h and to the resulting solution was added 1.0 M toluene solution of Et<sub>2</sub>AlCN (8.7 ml). The reaction was stirred at rt for 2 h, diluted with EtOAc (10 ml) and quenched with water (3 ml). The resulting slurry was then filtered through celite, concentrated and chromatographed to yield 1.1 g (74%) of (66).


**Step 6:** To a stirred solution of (66) (0.225 g, 0.398 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added TFA (1 ml). The mixture was stirred at rt for 1 h and concentrated. The residue was then diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with 10% NaOH. The extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to yield crude amine which was used directly in the next step without purification 0.164 g (89%).

To a stirred solution of crude amine (0.053 g, 0.115 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) was added Et<sub>3</sub>N (0.100 ml) followed by n-PrSO<sub>2</sub>Cl (0.05 ml) and stirred at rt for 1 h. The reaction was quenched with 2N NaOH and the organic layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined

-80-

organic layers were dried ( $\text{NaSO}_4$ ), filtered, concentrated and chromatographed to afford 0.025 g (50%) of (67).


- Step 7:** To a stirred solution of (67) (0.018 g, 0.031 mmol) in THF (2 ml) was added 3.0 M solution of  $\text{MeMgBr}$  in  $\text{Et}_2\text{O}$  dropwise at rt. The reaction was stirred for 2 h at rt and quenched with  $\text{H}_2\text{O}$ . The aqueous layer was then extracted with  $\text{EtOAc}$ , dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and chromatographed to yield 0.010 g (60%) of **126**. LRMS: calc'd: 562; found ( $\text{M}+\text{H}^+$ ) 563.

**Example 26**

10

Intermediate (1) (0.029 g) was dissolved in pyridine (0.1 ml) and isatoic anhydride (0.044 g) was added. The reaction solution was stirred for 48 h at room temperature then diluted with  $\text{EtOAc}$  (10 ml), washed with water, dried over  $\text{MgSO}_4$ , filtered and evaporated to give a solid.

- 15 The acetone-soluble portion of this solid was purified by thin layer chromatography (silica gel adsorbent; 95:5  $\text{EtOAc}:\text{Et}_3\text{N}$  eluant) to give a light colored foam (0.029 g) in 65% yield. MP (of hydrochloride): decomp  $>205^\circ\text{C}$

**Example 27**


20

- To a solution of **127** (0.096 g) and  $\text{CH}_2\text{Cl}_2$  (0.32 ml) was added pyridine (0.01 ml), acetic anhydride (0.02 ml) and N, N-dimethyl aminopyridine (4.8 mg). The resulting solution was stirred for 1.5 h, water (0.4 ml) was added and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3 X 1 ml). The organic extract was dried over  $\text{MgSO}_4$ , filtered and evaporated to give a solid residue which was purified by thin layer chromatography (silica gel adsorbent; 9:1  $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}$  eluant) to



25

-81-

give light colored foam (0.036 g) in 37% yield. MP (of hydrochloride): decomp  $>181^{\circ}\text{C}$ .

**Example 29**

- 5 To a solution of the acetate shown above (prepared as described in Example 1) (0.071 g), and  $\text{CH}_3\text{OH}$  (0.32 ml) was added  $\text{Et}_3\text{N}$  in three portions, over a 3 h period (0.06 ml total). The resulting solution was stirred 3 days at room temperature, water was added and the solution was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic extracts were dried over
- 10  $\text{MgSO}_4$ , filtered and evaporated to give the product (0.075 g) in 100% yield. M.p. (of hydrochloride): decomp  $>130^{\circ}\text{C}$ .

**Example 29**Step 1:



- 15 To a cooled ( $-78^{\circ}\text{C}$ ) solution of intermediate (3b) (0.41 g; 0.88 mmol) and THF (4.2 ml) was added  $n\text{BuLi}$  (0.34 ml of a 2.7 M solution in heptane) and the resulting solution was stirred for 3 min. Pyridine-3-uocarboxaldehyde was added as a solution in THF (0.17 ml in 2 ml, respectively) dropwise, over 1.5 min. The solution was stirred for 10 min
- 20 at low temperature, 40 min at room temperature, poured into water,

-82-

extracted with EtOAc, washed with brine and dried over  $\text{MgSO}_4$ . After filtration and evaporation, the resulting crude yellow oil was chromatographed on silica gel (77 g), eluting with 76:19:5 EtOAc:hexanes: $\text{Et}_3\text{N}$ . Evaporation of the appropriate fractions gave, as a diastereomeric mixture of carbinols, (37) as a clear oil (0.28 g) in 64% yield. HRMS:  $\text{MH}^+$ :  $\text{C}_{29}\text{H}_{43}\text{N}_4\text{O}_3$ : calc'd: 495.3335; found: 495.3319.

**Step 2:**

Treat the product of Step 1 with o-toluoyl chloride using a procedure similar to that described in Example 3 to obtain the desired compound **130**. MP (of hydrochloride): 174-177°C.

**Examples 30A and 30B****30A:****Steps 1 to 4:****Step 1:**

To a cooled (-78°C) solution of (24b) (17.9 g) and THF (170 ml) was added  $\text{CH}_3\text{Li}$  (43.2 ml of a 1.4 M solution in hexane) and the resulting solution was stirred for 10 min.  $\text{N-BuLi}$  (24.0 ml of a 2.5 M solution in hexane) was added and the resulting solution was stirred for 15 min. Piperonal was added as a solution in THF (9.6 g in 30 ml, respectively) dropwise and the cooling bath was removed. The solution was stirred for 15 min at room temperature, poured into dilute HCl and  $\text{Et}_2\text{O}$  and the  $\text{Et}_2\text{O}$  extracts were dried over  $\text{MgSO}_4$ . After filtration and evaporation, the resulting crude yellow oil was chromatographed on silica gel, eluting with 1:1 hexane: $\text{CH}_2\text{Cl}_2$  then 0→20 %  $\text{Et}_2\text{O}$ : $\text{CH}_2\text{Cl}_2$ .

-83-

The fractions containing the desired product were combined and evaporated to give as a mixture of carbinols (38) (16.1 g) in 75% yield.

Step 2:

To a solution of (38) (15.8 g),  $\text{CH}_2\text{Cl}_2$  (300 ml) and  $\text{Et}_3\text{SiH}$  (25 ml) was added TFA (25 ml). After 15 min at room temperature, the solution was washed with water, aqueous  $\text{NaHCO}_3$ , dried over  $\text{MgSO}_4$  and filtered through a pad (20g) of silica, washing through the product with  $\text{CH}_2\text{Cl}_2$ . Evaporation and trituration with hexane (200 ml) gave, after filtration and drying, a white solid, (39) (13.9 g) in 92% yield. M.p.: 99-101°C.


Step 3:

To a solution of (39) (2.23 g, 6.37 mmol),  $\text{CH}_2\text{Cl}_2$  (23 ml) and iodine (1.84 g) was added, in one portion, silver trifluoroacetate (1.60 g). The resulting mixture was stirred for 1.5 h at room temperature, stored at 5°C for 14 h, diluted with  $\text{CH}_2\text{Cl}_2$  (126 ml) and filtered. The resulting  $\text{CH}_2\text{Cl}_2$  extract was poured onto a column of silica gel (63 g) and the product washed through with  $\text{CH}_2\text{Cl}_2$ . Any fractions containing a light pink color were washed with 10%  $\text{Na}_2\text{S}_2\text{O}_3$  and dried, then combined with the other fractions and evaporated, to give (40) (2.86 g) as a white solid in 94% yield. M.p.: 125-127°C.

Step 4:

(40) (1.50 g, 3.14 mmol) was dissolved in N,N-dimethylacetamide (10 ml) and  $\text{CuCN}$  (0.94 g) and  $\text{NaI}$  (0.14 g) were added. The resulting mixture was degassed and heated at 110°C for 8 h. The mixture was cooled, diluted with  $\text{CH}_2\text{Cl}_2$ , filtered and the resulting solids were washed with  $\text{CH}_2\text{Cl}_2$ . The combined  $\text{CH}_2\text{Cl}_2$  extracts were washed with 1:1  $\text{NH}_4\text{OH}:\text{H}_2\text{O}$  (3X), water and brine, dried over  $\text{MgSO}_4$ , filtered, evaporated and applied to a silica gel column (55 g). After elution with hexanes (100 ml) and 4:1 hexanes:EtOAc, the desired fractions were evaporated to give a waxy white solid, (41), (1.10 g) in 94% yield. M.p.: 114-116°C.


-84-

Steps 5 and 6:Step 5:

- Treat (41) using a similar procedure to that described in Example 10, Step 3, to give the amine (42) as a white solid in 70% yield.  
M.p.: 55-58°C

Step 6:

- To a cooled (0°C) solution of diol (43) (0.88g), CH<sub>2</sub>Cl<sub>2</sub> (29 ml) and Et<sub>3</sub>N (1.21 ml) was added CH<sub>3</sub>SO<sub>2</sub>Cl (0.46 ml). The resulting solution was stirred at 0°C for 15 min and at room temperature for 1 h. The solution was poured into a mixture of CH<sub>2</sub>Cl<sub>2</sub> and ice water, the CH<sub>2</sub>Cl<sub>2</sub> layer was removed and washed with water, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration and careful evaporation of the solvent, CH<sub>3</sub>CN (2.30 ml) and iPr<sub>2</sub>EtN (1.20 ml) and (42) (0.64 g) were added and the mixture was heated at 80°C for 12 h. The resulting solution was diluted with EtOAc, washed with 2 N NaOH, water and brine, dried over MgSO<sub>4</sub>, filtered and evaporated to give a gold foam which was applied to a silica gel (91 g) column and eluted with 3:1 hexane:EtOAc, then 2:1 hexane:EtOAc. The desired fractions were combined and evaporated to give (44) (1.09 g) as a gold foam in 87% yield.  
HRMS:MH<sup>+</sup>: C<sub>28</sub>H<sub>29</sub>O<sub>6</sub>N<sub>4</sub>S: calc'd: 549.1808; measured: 549.1803.


Step 7:

131



-85-



- Part A: To a cooled (0°C) mixture of piperidone hydrate hydrochloride (27.5 g), CH<sub>2</sub>Cl<sub>2</sub> (350 ml), K<sub>2</sub>CO<sub>3</sub> (50 g) and water (250 ml) was added dropwise a solution of o-toluoyl chloride (22 ml) and CH<sub>2</sub>Cl<sub>2</sub> (25 ml). After the addition was complete, the cooling bath was removed and the mixture was stirred for 24 h, the CH<sub>2</sub>Cl<sub>2</sub> layer was removed, washed with water, dried over MgSO<sub>4</sub>, filtered and evaporated to give a clear oil. The oil was chromatographed on silica gel (320 g), eluting with 4:1 hexane:EtOAc and the desired fractions were combined and evaporated. The resulting clear oil crystallized on standing to give a white solid, (46), (35.7 g) in 92% yield.
- Part B: To a solution of (44) (0.21 g) and DMF (0.75 ml) was added PhSH (0.047 ml) and K<sub>2</sub>CO<sub>3</sub> (0.15 g) and the resulting mixture was stirred for 28 h. The reaction mixture was diluted with EtOAc and water and 2 N NaOH (1.0 ml) were added. The aqueous layer was extracted with EtOAc (three portions), the EtOAc extracts were combined and washed with brine, dried over MgSO<sub>4</sub>, filtered and evaporated to give 0.55 g of an oil (45) which was taken up in CH<sub>2</sub>Cl<sub>2</sub> (0.55 ml) and acetic acid (9 µl). To this solution was added (46) (0.04 g) and NaB(OAc)<sub>3</sub>H (0.048 g). After stirring for 6 h, the reaction solution was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 2 N NaOH. The CH<sub>2</sub>Cl<sub>2</sub> layer was washed with water and brine, dried over MgSO<sub>4</sub>, filtered, evaporated and applied to a thin layer silica plate. After elution with 2:1 CH<sub>2</sub>Cl<sub>2</sub>:acetone, the desired product, **131**, was collected as a foam (0.72 g) in 85% yield. M.p. (hydrochloride): 165-167°C.
- 30B:**

**132**

- To a solution of **131** (0.029 g) and THF (0.5 ml) was added nBu<sub>4</sub>NOH (0.085 ml) and 30% H<sub>2</sub>O<sub>2</sub> (0.017 ml). More nBu<sub>4</sub>NOH and H<sub>2</sub>O<sub>2</sub> were added as described above after 3 h and 48 h then the reaction was stirred 3 h more after the last addition. The reaction was diluted with EtOAc, the aqueous layer was removed, extracted with

-86-

EtOAc (3X) and the EtOAc extracts were combined, washed with water and brine, dried over  $\text{MgSO}_4$ , filtered and evaporated to give a white crystalline solid which was further purified by chromatography (silica gel; 100 mg; 95:5 EtOAc:Et<sub>3</sub>N eluant). The desired product was collected and evaporated to give a white solid, **132**, (10.5 mg) in 35% yield. M.p. (of hydrochloride): decomp >164°C.


**Example 31****Part A:**

To a solution of piperazine (13 g), N-BOC 4-piperidone (10 g),  $\text{CH}_2\text{Cl}_2$  (200 ml) and acetic acid (2.0 ml) was added  $\text{NaB}(\text{OAc})_3\text{H}$  (20 g) in four portions over 2 h. The mixture was stirred for 22 h, diluted with  $\text{CH}_2\text{Cl}_2$  and added slowly to 1 N NaOH (250 ml). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3X), the  $\text{CH}_2\text{Cl}_2$  extracts were combined, washed with water and dried over  $\text{MgSO}_4$ . The solution was filtered, concentrated and purified by silica gel chromatography (eluant: 8:1  $\text{CH}_2\text{Cl}_2$  :  $\text{CH}_3\text{OH}$ , then 10:5:1  $\text{CH}_2\text{Cl}_2$  :  $\text{CH}_3\text{OH}$  :  $\text{NH}_3(\text{aq})$ ). After evaporation of the solvent, (47) was collected as a white solid (7.18 g) in 53 % yield.

To a solution of p-bromo benzaldehyde (5.25 g),  $\text{CH}_2\text{Cl}_2$  (100 ml) and (47) (8.72 g) was added  $\text{NaB}(\text{OAc})_3\text{H}$  (8.05 g) in 8 portions over 40 min. After stirring for 19 h,  $\text{CH}_2\text{Cl}_2$  (20 ml) and  $\text{NaB}(\text{OAc})_3\text{H}$  (1.05 g) were added, the reaction mixture was stirred for 2 h, diluted with  $\text{CH}_2\text{Cl}_2$ , poured into 2 N NaOH and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$ . The  $\text{CH}_2\text{Cl}_2$  extracts were washed with water and brine, dried over  $\text{MgSO}_4$ , filtered, evaporated and chromatographed on silica gel (250 g), eluting with 76:19:5 hexane:EtOAc:Et<sub>3</sub>N. The desired fractions



-87-

were combined and evaporated to give (48) as a white solid (8.48 g) in 69 % yield. M.p.: 87-90°C.

**Part B:**

- 5 To a solution of m-toluene sulfonyl chloride (13.2 g) and acetone (107 ml) was added a solution of KF (11 g) and water (87 ml), slowly dropwise. The resulting solution was diluted with EtOAc and the organic layer was removed. The aqueous layer was extracted with EtOAc, combined with the initial EtOAc extract, dried over MgSO<sub>4</sub>, filtered and
- 10 evaporated to give (49), (11.7 g) in 97% yield, which was used directly without further purification.

- (50) was prepared using a procedure similar to Example 29, except (49) was used in place of 3-pyridine carboxaldehyde. Using procedures described in Example 1 (or Example 2), (50) was converted
- 15 to **133**. M.p. (hydrochloride): decomp >227°C.

**Example 32****Part A:**

- 20 The amino acid (0.21 g), DME (15 ml) and 1 M NaOH (3 ml) were mixed and allyl chloroformate (0.2 ml) was added. The solution was stirred for 15 h, cooled to 0°C, acidified with 1 M HCl and extracted with EtOAc. The EtOAc extracts were washed with brine and dried over

-88-


Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give (51) as a colorless foam, which was used directly in the next procedure.

Cyanuric fluoride (0.41 g), (51) (0.50 g), pyridine (0.15 ml) and CH<sub>2</sub>Cl<sub>2</sub> (15 ml) were mixed at 0°C. The mixture was stirred at 0°C for 3.75 h, diluted with CH<sub>2</sub>Cl<sub>2</sub> and poured into cold water. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the combined CH<sub>2</sub>Cl<sub>2</sub> extracts were dried over MgSO<sub>4</sub>, filtered and concentrated to give crude (52) (0.50 g) as a thick oil, which was used directly in the next procedure.


The acid fluoride (52) (0.31 g) and CH<sub>2</sub>Cl<sub>2</sub> (10 ml) were mixed and NaBH<sub>4</sub> (72 mg) was added followed by CH<sub>3</sub>OH (0.9 ml), dropwise over 10 min. The reaction was stirred for 20 min at room temperature, poured into a separatory funnel containing 1 M HCl and EtOAc and the aqueous layer extracted with EtOAc. The organic extracts were combined, washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to give (53) as an oil (0.30 g) in 51 % yield.

Palladium acetate (4 mg), 3,3',3''-phosphinidynetris(benzene sulfonic acid) sodium salt (20 mg), Et<sub>2</sub>NH (1.3 g) and (53) (0.28 g) were mixed with CH<sub>3</sub>CN (3 ml) and water (3 ml) and stirred for 3 h. The reaction mixture was co-evaporated with toluene (6 ml) to obtain (54), which was used directly in the next part.

#### Part B:




-89-



- (55) (0.50 g) (prepared using the procedure of Example 1, using p-fluorobenzaldehyde) was taken up in CH<sub>3</sub>OH (20 ml) and CH<sub>2</sub>Cl<sub>2</sub> (10 ml), cooled to 0°C and NaBH<sub>4</sub> (80 mg) was added. After stirring for 2.5 h
- 5 at room temperature, the solution was partitioned between EtOAc and 1 M HCl. The EtOAc layer was removed, the aqueous layer was extracted with EtOAc, the organic extracts were combined, washed with brine, dried over MgSO<sub>4</sub>, filtered and evaporated to give (56) (0.52 g) as a colorless oil. A solution of CH<sub>2</sub>Cl<sub>2</sub>, SOCl<sub>2</sub> (0.13 ml) and DMF (0.033 ml)
- 10 was stirred for 4 h at room temperature and concentrated. The residue was co-evaporated with toluene (2x 20 ml) to give a yellow solid (57) (0.20 g) which was dissolved in CH<sub>3</sub>CN (10 ml) and diethanolamine (0.24 g). After heating at reflux for 2 h, the reaction solution was
- 15 partitioned between water and EtOAc, the EtOAc layer was removed, the aqueous layer was extracted with EtOAc, the organic extracts were combined, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated to give the product (58) (0.26 g) as a colorless oil. This product was dissolved in 1,2-dichloroethane (5 ml), SOCl<sub>2</sub> (0.16 ml) was added and the resulting mixture was heated at reflux for 3 h. The volatile
- 20 materials were removed in vacuo to give the product (59) as a thick gum, which solidified on storage at 5°C. This material (0.17 g) was mixed with (54) (0.10 g), NaI (0.20 g), CH<sub>3</sub>CN (10 ml) and diisopropyl ethyl amine (0.45 ml) and the resulting mixture was heated at reflux for 3.5 h. The reaction solution was partitioned between water and EtOAc, the EtOAc
- 25 layer was removed, the aqueous layer was extracted with EtOAc, the organic extracts were combined, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated to give a brown oil. Purification on silica gel, eluting with 10:1 CH<sub>2</sub>Cl<sub>2</sub> :CH<sub>3</sub>OH, gave (60) (77 mg) as an oil. HRMS:MH<sup>+</sup>:C<sub>29</sub>H<sub>40</sub>N<sub>3</sub>O<sub>7</sub>S calc'd: 574.2587; measured: 574.2580.
- 30 (60) (77 mg) was dissolved in 1 M HCl in CH<sub>3</sub>OH (5 ml), stirred for 24 h, concentrated and the residue was taken up in CH<sub>3</sub>OH and

-90-

trituated with Et<sub>2</sub>O to give a white solid (59 mg). Using the procedure of Example 1, (60) was converted to **134**. HRMS (FAB) calc'd for C<sub>32</sub>H<sub>38</sub>N<sub>3</sub>O<sub>6</sub>S (MH<sup>+</sup>): 592.2481; found: 592.2467; m.p. 187-197°C.

**Example 33**

**Step 1:** To a cooled (-78°C) mixture of 4-amino benzyl alcohol (0.50 g), diisopropyl ethyl amine (1.41 ml) and CH<sub>2</sub>Cl<sub>2</sub> (40 ml) was added TMSCl (0.52 ml). The resulting slurry was stirred at -78°C for 15 min and at 0°C for 1 h. The resulting solution was cooled to -78°C, p-methoxyphenyl sulfonyl chloride (0.84 g) was added, the cooling bath was removed and the reaction was stirred at room temperature for 15 h. The reaction solution was diluted with CH<sub>2</sub>Cl<sub>2</sub> and water was added. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>, the combined organic extracts were dried over MgSO<sub>4</sub>, filtered and evaporated to give a yellow oil, which was taken up in CH<sub>3</sub>OH (20 ml) and K<sub>2</sub>CO<sub>3</sub> (1.5 g) was added. The resulting mixture was stirred for 1 h., filtered through a plug of Celite, concentrated, taken up in EtOAc and washed with water. The aqueous layer was extracted with EtOAc, the combined organic extracts were dried over MgSO<sub>4</sub>, filtered and evaporated to give a yellow oil (0.78 g), which was used without further purification in the next step.

-91-


**Step 2:** (68) (0.47 g) and  $\text{MnO}_2$  (3.0 g) were mixed with  $\text{CH}_2\text{Cl}_2$  (20 ml) and stirred for 20 h. The solution was filtered through Celite and concentrated to give 0.30 g of a thick oil, (69), which was used directly in the next step.

- 5 **Step 3:** Use a procedure similar to the second portion of Part A in Example 31, with intermediates (69) and (70) as the reactants. Crude (71) (0.30 g) was used directly in the next step.

10 **Step 4:** Treat (71) (148 mg) using a procedure similar to that in the last portion of Part A of Example 32 to obtain the corresponding brown crude amine, which was used directly in the next step.

**Step 4:** Using the procedure of in Example 1, treat (71) to obtain **135**. M.p. (with decomposition) 190-200°C.

### Example 34



15

**Step 1:** To a stirred 0°C solution of 4,4'-bispiperidine hydrochloride (2.53 g, 10.5 mmol) in a 1:1 solution of  $\text{Et}_2\text{O}$ /10% aq.  $\text{NaOH}$  (26 ml) was added a 2.5 M solution of di-*tert*-butyldicarbonate in  $\text{Et}_2\text{O}$  (10 ml, 23.1 mmol) over 30 min with vigorous stirring. After stirring at room

-92-

temperature for 2 h, the mixture was poured into a separatory funnel, extracted with Et<sub>2</sub>O, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The resulting product (3.00 g) was used without further purification.

5           The crude bispiperidine (2.74 g, 7.44 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 ml) and TFA (0.57 ml, 7.44 mmol) was added to the solution. The reaction stirred at room temperature and monitored by TLC. After two more equivalents of TFA were added, the reaction was approximately a 1:2:1 mixture of starting material, (72), and 4,4'-  
10 bispiperidine. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with 1N NaOH, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography (10/90 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>, then 5/10/85 NH<sub>4</sub>OH/CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) yielded 1.00 g of (72) (53% yield).

Step 2: To a solution of (73) (480 mg, 1.85 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7.4 ml)  
15 was added (72) (520 mg, 2.04 mmol) and sodium triacetoxymethylborohydride (590 mg, 2.78 mmol). The reaction was stirred for 12 h at room temperature under N<sub>2</sub>. The mixture was quenched with NaHCO<sub>3</sub> (sat), extracted with EtOAc, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure, yielding (74) (880 mg, 94% yield).

20 Step 3: To a solution of (74) (880 mg, 1.72 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8.6 ml) was added CH<sub>3</sub>SO<sub>3</sub>H (0.17 ml, 2.58 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4.7 ml). After cooling to 0°C with stirring, 60% MCPBA (1.04 g, 3.61 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added dropwise. The reaction was stirred for 1 h at room  
25 temperature under N<sub>2</sub>. The mixture was washed with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (sat), 1 N NaOH, and H<sub>2</sub>O, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield 0.93 g of the sulfone (100% yield).

The sulfone (0.93 g, 1.71 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (11.4 ml) and TFA (2.6 ml, 34.2 mmol) was added. The reaction was stirred for 2 h at room temperature under N<sub>2</sub>. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>,  
30 washed with 1N NaOH, filtered, and concentrated under reduced pressure, yielding the free amine (0.45 g, 59% yield).


To a solution of the amine (23.6 mg, 53.3 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.53 ml) was added *o*-toluoyl chloride (8.3 μL, 64 μmol) and Et<sub>3</sub>N (11 μL, 80 μmol). The reaction was stirred for 2 h at room temperature under N<sub>2</sub>.  
35 The mixture was concentrated under reduced pressure and purified by



-93-

PTLC (10/90 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>), concentrated in the presence of 1N HCl/Et<sub>2</sub>O, yielding **136** (15.8 mg, 50% over three steps).


LRMS calc'd: 560; found (M+H): 561.

**Example 35**

- NaH (0.59 g, 14.6 mmol, 60% purity) was suspended in DMF (10 ml) followed by addition of 2-propanethiol (1 ml, 11.2 mmol) at 0°C under N<sub>2</sub>. The reaction mixture was stirred at room temperature for 5 min. (75) (2.94 g, 7.5 mmol) in DMF (10 ml) was added dropwise. The reaction mixture was heated with stirring for 2 hrs. The reaction mixture was then diluted with 1N NaOH (40 ml) and extracted with Et<sub>2</sub>O (3x 80 ml). The organic extract was dried with NaHCO<sub>3</sub>. Removal of solvent and recrystallization with Et<sub>2</sub>O afforded (76) (1.08 g, 32%).
- (76) (1.08 g, 2.4 mmol) was dissolved in CH<sub>3</sub>OH (10 ml) followed by addition of NaBH<sub>4</sub> (0.14 g, 3.6 mmol). The reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (200 ml) followed by washing with 1N NaOH (50 ml). The organic solution was dried with NaHCO<sub>3</sub> and concentrated to give solid residue which was then dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 ml). To this solution was added Et<sub>3</sub>SiH (1.5 ml) followed by TFA (3 ml). The reaction mixture was stirred at room temperature for 1hr. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (100 ml), washed with 1N NaOH (100 ml), and dried with NaHCO<sub>3</sub>. Removal of solvent afforded pure (77) (0.7 g, 88%).
- (77) (49 mg, 0.15 mmol) was dissolved in a solution of CH<sub>2</sub>Cl<sub>2</sub> (3 ml) and Et<sub>3</sub>N (0.3 ml) followed by addition of 1-naphthoyl chloride (0.5 ml). The reaction mixture was stirred for 30 min and separated by PTLC

-94-

to afford **137** (53 mg, 74% ) as free base. HRMS: calc'd: (M+1): 487.2783; found 487.2798.

**Example 36**

5 (78) (1 g, 3.6 mmol) was dissolved in a solution of  $\text{CH}_2\text{Cl}_2$  (20 ml) and  $\text{Et}_3\text{N}$  (1.5 ml) followed by addition of  $(\text{CF}_3\text{CO})_2\text{O}$  (0.82 ml, 5.7 mmol). The reaction mixture was stirred at room temperature overnight. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (150 ml), washed with 1N HCl (50 ml) followed by 1N NaOH (50 ml), and dried with  $\text{MgSO}_4$ .  
10 Removal of solvent afforded crude (79).

(79) was dissolved in  $\text{CH}_3\text{OH}$  (25 ml) followed by addition of  $\text{NaBH}_4$  (0.2 g, 5.3 mmol). The reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (100 ml) followed by washing with 1N NaOH (50 ml). The organic solution was dried with  $\text{MgSO}_4$  and concentrated to give a residue which was then dissolved in  $\text{CH}_2\text{Cl}_2$  (30 ml). To this solution was added  $\text{Et}_3\text{SiH}$  (3 ml) followed by TFA (10 ml). The reaction mixture was stirred at room temperature for 4 hrs. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (200 ml), washed with 1N NaOH (100 ml), and dried with  $\text{MgSO}_4$ .  
20 Removed solvent to afford crude (80).

(80) was dissolved in  $\text{CH}_2\text{Cl}_2$  (5 ml) followed by addition of 1M  $\text{BBr}_3/\text{CH}_2\text{Cl}_2$  (8.6 ml) at  $0^\circ\text{C}$ . The reaction mixture was stirred at  $0^\circ\text{C}$  for

-95-


1 hr, diluted with  $\text{CH}_2\text{Cl}_2$  (100 ml), washed with 10%  $\text{NaHCO}_3$ , and dried with  $\text{MgSO}_4$ . Removed solvent followed by column chromatography (20% EtOAc/hexane) to afford (81) (0.725 g, 58% from (78)).

5 (81) (0.35 g, 1.2 mmol), 2-propanol (0.084 g, 1.4 mmol) and  $\text{PPh}_3$  (0.38 g, 1.4 mmol) were dissolved in THF (5 ml) under  $\text{N}_2$ . The reaction mixture was cooled to  $0^\circ\text{C}$  followed by addition of DEAD (0.27 ml, 1.1 mmol). The reaction mixture was stirred at room temperature overnight and separated by PTLC without workup to afford (82) (0.3 g, 75 %).

10 (82) (0.3 g, 0.9 mmol) was dissolved in a solution of  $\text{CH}_3\text{OH}/\text{water}$  (5:1) (10 ml) followed by addition of  $\text{K}_2\text{CO}_3$  (0.5 g, 3.6 mmol). The reaction mixture was stirred at room temperature for 1h. Standard aqueous workup afforded (83) (0.203 g, 96%).

(83) was converted to 138 following the procedures described in  
15 Example 35. HRMS: calc'd: (M+1): 471.3012; found 471.3009.

### Example 37



20 (84) (0.85 g, 3.4 mmol) and N-BOC-4-piperidone (1 g, 5.0 mmol) were dissolved in 1,2-dichloroethane (50 ml) followed by addition of  $\text{Na}(\text{OAc})_3\text{BH}$  (2.2 g, 10.1 mmol). The reaction mixture was stirred at room temperature for two days. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (200 ml), washed with 1N  $\text{NaOH}$  (100 ml) and dried with

-96-


NaHCO<sub>3</sub>. Column chromatography (5% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) afforded (85) (1.27 g, 87%).

(85) (0.2 g, 0.46 mmol) was dissolved in THF (8 ml) under N<sub>2</sub> and cooled down to -78°C followed by addition of 2.5 M n-BuLi (0.3 ml, 0.73 mmol). The reaction mixture was stirred at -78°C for 45 min followed by addition of a solution of (86) (0.06 g, 0.83 mmol) in THF (1 ml). The reaction mixture was stirred at -78°C for 30 min, diluted with CH<sub>2</sub>Cl<sub>2</sub> (100ml), washed with water (50 ml), and dried with NaHCO<sub>3</sub>. The organic solution was concentrated and column chromatography (5% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) afforded (87) (0.135 g, 69%).

To a mixture of (87) (0.135 g, 0.32 mmol) and Et<sub>3</sub>SiH (1ml) was added 20% TFA/CH<sub>2</sub>Cl<sub>2</sub> (5 ml). The reaction mixture was stirred at room temperature for 1h and concentrated to give (88) (0.09 g, 91%).

(88) was treated in a manner similar to that described in Example 35 to obtain 139. HRMS: calc'd: (M+1): 467.3062; found 467.3066.

#### Examples 38A and 38B



To a suspension of 500 mg of NaH (60% in mineral oil) in 60 ml of DMF was added slowly 1.0 g of tert-butyl thiol. The mixture was stirred at room temperature for 30 min. To this mixture, 3.1 g (10 mmol) of ketone

-97-

(89) was added dropwise. The mixture was heated at 70°C overnight. After cooling down to room temperature, the mixture was quenched with 100 ml of water. The aqueous phase was extracted with 3X100 ml of CH<sub>2</sub>Cl<sub>2</sub>. The combined organic phases were dried over MgSO<sub>4</sub>, filtered and concentrated. The crude (90) was purified by column chromatography on silica gel using 5% EtOAc-Hexanes (2.65 g, 69%).

5 To a solution of 2.65 g of (90) in 40 ml of CH<sub>2</sub>Cl<sub>2</sub> was added 10 ml of TFA. The mixture was stirred at room temperature for 2 hours and then was concentrated. The residue was taken up in 100 ml of CH<sub>2</sub>Cl<sub>2</sub>,  
10 washed with 10% NaOH. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give a crude (91) (1.8 g, 92%).


A mixture of (91) (1.8 g), N-Boc-4-piperidone (1.5 g, 1.1 eq.), NaB(OAc)<sub>3</sub>H (2.8 g, 2 eq.) and 0.5 ml HOAc in 40 ml of 1,2-dichloroethane was stirred at room temperature over night. The mixture was  
15 quenched with 10% NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude (92) was purified by column chromatography on silica gel using 20% EtOH/EtOAc (2.7 g, 90.3%).

To a solution of (92) (2.5 g) in 100 ml of CH<sub>3</sub>OH was added 500  
20 mg (2 eq.) of NaBH<sub>4</sub>. The mixture was stirred at room temperature for 1 hour and then was concentrated to dryness. The residue was taken up in 70 ml of 10% NaOH and extracted with 3X70 ml of CH<sub>2</sub>Cl<sub>2</sub>. The combined organic phases were dried over MgSO<sub>4</sub>, filtered and concentrated (1.8 g crude). This crude was dissolved in 40 ml of CH<sub>2</sub>Cl<sub>2</sub>  
25 and 1ml of Et<sub>3</sub>SiH. To this mixture was added 20 ml of TFA and stirred at room temperature for 1 hour. The mixture was concentrated to dryness and the residue was washed with cold hexanes. The residue was taken up in 100 ml of CH<sub>2</sub>Cl<sub>2</sub> and washed with 10% NaOH. The combined organic phases were dried over MgSO<sub>4</sub>, filtered and  
30 concentrated to give a crude (93) (810 mg, 43%).

(93) and 1.3 ml (4 eq.) of Et<sub>3</sub>N were dissolved in 40 ml of 1,2-dichloroethane and this solution was used for parallel synthesis to react with different acid chlorides. Typical yield of the coupling reaction is 80%. The HCl salts were prepared by adding HCl·OEt<sub>2</sub> to the free base.  
35 140: LRMS calc'd: 500; found 501; 141: LRMS calc'd: 478; found 479.

-98-

## Example 39



5 A mixture of (89) (7.9 g, 25.6 mmol), sesamol (1.1 g, 1 eq.) and  $\text{K}_2\text{CO}_3$  (1.5 g) in 40 ml of dimethylacetamide was heated at 150°C over night. The mixture was allowed to cool to room temperature and then was quenched with 200 ml of water. The aqueous phase was extracted with 3X150 ml of  $\text{CH}_2\text{Cl}_2$ . The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude (94) was purified by column chromatography on silica gel using 20% EtOAc-Hexanes (2.8 g, 26%).


15 To a solution of (94) (2.8 g) dissolved in 50 ml of  $\text{CH}_2\text{Cl}_2$  was added 5 ml of TFA. The mixture was stirred at room temperature for 1 h and was then concentrated to dryness. The residue was taken up in 150 ml of  $\text{CH}_2\text{Cl}_2$ , washed with 100 ml of 10% NaOH. The aqueous layer was extracted with 3X100 ml of  $\text{CH}_2\text{Cl}_2$ . The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and concentrated to give crude (95) as a solid (1.7 g, 79%).

20 The crude (95) (1.7 g, 5.2 mmol) was treated with N-BOC-4-piperidone,  $\text{NaB(OAc)}_3\text{H}$  and acetic acid (1.3 g, 4 eq.) in a manner similar to that described in Example 38 to obtain crude (96) (2.3 g, 87%).

-99-

The BOC-group was removed to give a crude (97) (1.7 g, 93%), which was reacted with 2-toluoyl chloride as described in Example 14, Step 8, followed by purification by TLC on silica gel using 20% EtOH-EtOAc to obtain **142** (197 mg, 90%). The hydrochloride salt was prepared by adding HCl•Et<sub>2</sub>O to the free base. LRMS calc'd: 526; found: 527.

## Examples 40A and 40B




To a stirred solution of 500 mg of (98) in 10 ml of THF was added 0.6 ml of BuLi (2 M in hexane) at -78°C. The mixture was stirred at this temperature for 15 mins. and then a solution of 179 mg of the isocyanate in 15 ml of THF was added. The mixture was stirred at -78°C for 1 hour. The mixture was quenched with water, extracted with EtOAc. The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude (99) was purified by prep.TLC on silica gel using 25%EtOH-EtOAc (388 mg).



To a stirred solution of 328 mg of (99) in 50 ml of THF was added 41 mg (60% in mineral oil) NaH, followed by 85 mg of CH<sub>3</sub>I. The mixture was stirred at room temperature for 1 hour and then was purified by prep. TLC on silica gel using 25%EtOH-EtOAc to obtain 141 mg of **143**. LRMS calc'd: 593; found: 594.

-100-

To 92 mg of (99) in 20 ml of CH<sub>2</sub>Cl<sub>2</sub> was added 5 ml of TFA. The solution was stirred at RT for 1 hour, quenched with 10% NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (K<sub>2</sub>CO<sub>3</sub>), filtered and concentrated (66 mg crude). This crude was dissolved in 20 ml of CH<sub>2</sub>Cl<sub>2</sub>, added 0.25 ml of Et<sub>3</sub>N followed by 0.25 ml (1M in CH<sub>2</sub>Cl<sub>2</sub>) of 2,3-dimethylbenzoyl chloride. The mixture was stirred at RT for 1 hour and was purified by prep. TLC using 25%EtOH-EtOAc to obtain 81 mg **144**. LRMS calc'd: 611; found: 612.

**Example 41****145**


To a stirred solution of **83** (255 mg) in 50 ml of CH<sub>3</sub>OH was added 186 mg of NaBH<sub>4</sub>, the mixture was stirred at room temperature for 30 min. and was then quenched water. The mixture was extracted with EtOAc and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give a yellowish oil (286 mg). The crude was dissolved in 10 ml of CH<sub>2</sub>Cl<sub>2</sub> and 2ml of Et<sub>3</sub>SiH. To this mixture, 2ml of TFA was added and the mixture was stirred at room temperature for 5 hours. The mixture was neutralized with 3N NaOH, the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude was purified by prep. TLC on silica gel using 25%EtOH-EtOAc (20 mg, 10%). LRMS: calc'd: 510; found: 511.

**Example 42****146**

25





-101-



To a stirred solution of (100) in 10 ml of DMSO was added 10 ml of Ac<sub>2</sub>O, the mixture was stirred at room temperature for 2 h, quenched with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude (101) was purified by prep. TLC on silica gel using 25%EtOH-EtOAc (110 mg). (101) was dissolved in 40 ml of CH<sub>2</sub>Cl<sub>2</sub>, and 5 ml of TFA was added. The mixture was stirred at room temperature for 30 min., concentrated to dryness. The residue was taken up in 100 ml of CH<sub>2</sub>Cl<sub>2</sub> and was washed with 10% NaOH. The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give a crude (102) (108 mg).

To a stirred solution of 28 mg of (102) and 0.1 ml of Et<sub>3</sub>N in 3 ml of CH<sub>2</sub>Cl<sub>2</sub> was added 0.25 ml of 2,3-dimethylbenzoyl chloride (1.0 M solution in CH<sub>2</sub>Cl<sub>2</sub>). The mixture was stirred at room temperature for 1.5 h. The crude was purified by prep. TLC on silica gel using 25%EtOH-EtOAc to obtain 146 (25 mg). LRMS calc'd: 596; found: 597.

**Example 43****Step1:**

20

m-Thiocresol (5.0 g), 1-bromo-4-iodobenzene (10.5 g), CuI (7.7 g) and K<sub>2</sub>CO<sub>3</sub> (20.0 g) in anhyd. DMF (100 ml) was heated at 130°C under N<sub>2</sub> overnight. The mixture was diluted with EtOAc and 10% NH<sub>4</sub>OH/NH<sub>4</sub>Cl and stirred, open to air, for 2 hours. The layers were separated, the organic phase was washed with the buffer solution 2x, followed by a water wash (3x). The organics were dried over sodium


25

-102-

sulfate, filtered and concentrated to a dark brown oil. The crude was flash chromatographed (silica gel) eluted with hexane to give a pale yellow oil which solidified on standing under vacuum (10.9g of (103)).

Step 2:

5




10

To a cold (-78°C) solution of (103) (2.0g) in anhyd. THF (50 ml) was slowly added n-BuLi (3.2 ml, 7.92 mmol) and the mixture was stirred at -78°C for 15 min. Ethyltrifluoroacetate (1.53 g, neat) was added in one portion and this mixture was stirred at -78°C to room temp over 3 h. The mixture was poured into water, THF was removed in vacuo and the aqueous residue was extracted with EtOAc. The organics were washed with brine, dried over anhyd. Na<sub>2</sub>SO<sub>4</sub> and concentrated to 1.97 g of a yellow oil, 2 (104)

Step 3:

15




20

To a solution of crude (104) (1.9g) and t-butyl -1-piperazine-carboxylate (3.6 g) in dry dichloroethane (20 ml) was added titanium IV chloride (6.4 ml, 6.4 mmol) in portions. This was stirred at room temp. for 2 days. The reaction was cooled to 0°C and NaBH<sub>3</sub>CN (1.2 g) in CH<sub>3</sub>OH (20 ml) was added. The mixture was stirred overnight at room temp, then filtered through a pad of celite. The layers were separated and the organic was washed with water and brine, then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to a pale red viscous oil. The crude was purified by column chromatography eluted with hexane (neat) to 10% EtOAc/hexane to give 1.7g of viscous pale yellow oil, (105).


25

Step 4:




-103-


To a solution of (105) (1.72 g) in  $\text{CH}_2\text{Cl}_2$  (25 ml) was added TFA (20 ml). The reaction was stirred for 1 hour, cooled to  $0^\circ\text{C}$  and neutralized with  $\text{NaHCO}_3$ . The aqueous phase was saturated with NaCl crystals and extracted with  $\text{CH}_2\text{Cl}_2$ . The organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to 1.32g of viscous pale yellow oil, (106).

**Step 5:**

A solution of (106) (1.32g) and 1-*t*-butyloxycarbonyl-4-piperidone (858 mg) in dichloroethane (15 ml) was treated with  $\text{NaBH}(\text{OAc})_3$  (1.1 g) followed by HOAc (0.42 ml, 7.4 mmol). The mixture was stirred at room temp under  $\text{N}_2$  for 4 days. The reaction was diluted with EtOAc, washed with 1N NaOH (2x), water, and brine. The organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The crude was purified by column chromatography eluted with 25% EtOAc/hexane to give 1.1g of viscous clear oil (107).

**Step 6:**

A solution of (107) (1.0 g) in anhyd.  $\text{CH}_2\text{Cl}_2$  (10 ml) was treated with TFA (10 ml) and stirred at room temp, under  $\text{N}_2$  for 1 hour. The mixture was then cooled to  $0^\circ\text{C}$ , neutralized with saturated  $\text{NaHCO}_3$  and extracted with  $\text{CH}_2\text{Cl}_2$ . The organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to 949 mg of pale yellow foam, (108).

**Step 7:**


To a solution of (108) (100 g) in anhyd.  $\text{CH}_2\text{Cl}_2$  (1.0 ml) containing  $\text{Et}_3\text{N}$  (38  $\mu\text{l}$ , 0.27 mmol) was added 1-naphthoyl chloride (41  $\mu\text{l}$ , 0.27 mmol). This solution was stirred overnight at room temp. The

-104-

crude was purified by directly applying to a Preparative TLC plate (1x2000 micron) eluted with 5% Et<sub>3</sub>N/EtOAc to give 86.4 mg of (109).  
Step 8:

To a solution of (109) (63.3mg) in anhyd. CH<sub>2</sub>Cl<sub>2</sub> (4.0 ml) was added 0.5M CH<sub>3</sub>SO<sub>2</sub>H solution in CH<sub>2</sub>Cl<sub>2</sub> (1.2 ml, 0.6 mmol). After stirring for 0.5 h, 30% H<sub>2</sub>O<sub>2</sub> (31μl, 0.6 mmol) was added and stirred overnight at room temp. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, neutralized with saturated NaHCO<sub>3</sub>. The layers were separated, dried over Na<sub>2</sub>SO<sub>4</sub> and applied to a Preparative TLC plate (2000 micron) eluted with 5% Et<sub>3</sub>N/EtOAc. Isolated 19.4 mg of **147**. HRMS calc'd: 636.2508; found: 636.2509.

#### Example 44



-105-

To a stirred solution of (110) (9.3 g, 53.2 mmol) and  $\text{NH}_4\text{NO}_3$  (4.4 g, 55.8 mmol) was added trifluoroacetic anhydride (40 ml) at  $-15^\circ\text{C}$ . The solvent was evaporated under vacuum at rt. The resulting oil was then diluted with water and the compound was extracted with  $\text{CH}_2\text{Cl}_2$  (2x50 ml). The combined organic layers were washed sequentially with dilute  $\text{NaHCO}_3$  and brine, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and chromatographed to afford (111) (3.8 g, 43%) as colorless oil.

To a stirred solution of (111) (3.8 g, 12.0 mmol) in  $\text{CH}_3\text{OH}:\text{H}_2\text{O}$  (5:1) 100 ml was added  $\text{K}_2\text{CO}_3$  (3.3 g, 24.0 mmol) and stirred for 3 h at rt. The reaction mixture was concentrated and poured into 2N NaOH and extracted with  $\text{CH}_2\text{Cl}_2$  (2x50 ml). The combined layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated to yield (112) (1.79 g, 70%) as a colorless oil.


(112) was treated with N-BOC-4-piperidone, then extracted as described in Example 38 to yield (113) (0.700g, 20%) as a colorless oil.

The BOC group on (113) was removed with TFA to obtain the amine (114), and (114) was reacted with 1-naphthoyl chloride as described in Example 14, step 8, to obtain (115) (0.448 g, 73%) as a colorless oil.


To a solution of (115) (0.170 g, 0.345 mmol) in  $\text{CH}_3\text{OH}$  (5 ml) was added  $\text{SnCl}_2$  (0.262 g, 1.38 mmol) and concentrated HCl (1 ml) and the reaction was heated to reflux for 12 h. The reaction mixture was then cooled to rt and poured into 10% NaOH and extracted with  $\text{CH}_2\text{Cl}_2$  (2x10 ml). The extracts were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated to yield (116) (0.150 g, 100%).

To a solution of (116) (0.02 g, 0.047 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml) was added  $\text{Et}_3\text{N}$  (0.013 ml, 0.094 mmol) followed by isobutyryl chloride (0.010 ml, 0.07 mmol) and the mixture stirred at rt for 2 h. The reaction mixture was then poured into sat'd aq  $\text{NaHCO}_3$  and extracted with  $\text{CH}_2\text{Cl}_2$  (2x5 ml). The extracts were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and chromatographed to yield 148 (0.014 g, 65%). LRMS: calc'd: 497; found ( $\text{M}+\text{H}^+$ ): 498.

-106-

**Example 45**

To a solution of (1) (20.0 mg, 0.042 mmol) in anhydrous THF (0.4 ml) at 0°C were successively added *n*-butyl sulfamoyl chloride (117) (literature prep.) (14.4 mg, 0.084 mmol) and Et<sub>3</sub>N (11.8 μl, 0.084 mmol) and the resulting mixture was stirred 14 h at room temperature. After evaporation of the solvent, the residue was treated with saturated aqueous NaHCO<sub>3</sub> solution (10 ml), extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 5 ml), dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. Purification of the crude by preparative silica gel chromatography (eluent CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>3</sub> 96:4:1) provided **149** (17.3 mg, 66%) as an oil: MH<sup>+</sup> = 621 (hydrochloride salt).

**Example 46**

-107-

To a stirred 0°C solution of (118) (50.0 g, 205 mmol) in a 1:1 solution of Et<sub>2</sub>O/10% aq. NaOH (500 ml) was added a 2.5 M solution of di-*tert*-butyldicarbonate in Et<sub>2</sub>O (100 ml, 246 mmol) over 30 min with vigorous stirring. After stirring at room temperature for 2 h, the mixture  
5 was extracted with Et<sub>2</sub>O, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The resulting product (73.9 g) was used without further purification.


3,4-Methylenedioxythiophenol (19.1 g, 124 mmol) was added dropwise to a stirred 0°C suspension of NaH (5.46 g of a 60%  
10 dispersion, 136 mmol) in DMF (75 ml). After stirring 30 min at 0°C, (118) (31.8 g, 103 mmol) in DMF (75 ml) was added and the solution warmed to room temperature overnight. The reaction was quenched with water and extracted with EtOAc. The organics were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The crude product was  
15 recrystallized from EtOAc/hexanes, with the first two crops yielding 15.8 g pure sulfide (35% yield).

The sulfide was treated with TFA to obtain the free amine, then reacted with N-BOC-4-piperidone as described in Example 38 to obtain  
(119) (8.20 g, 69% yield).


20 The BOC-group was removed and the resultant amine was reacted with 2-toluoyl chloride in a manner similar to that described in Example 14, Step 8. The mixture was concentrated under reduced pressure, yielding the aryl amide (0.65 g, 100% yield).

A solution of the aryl amide (0.21 g, 0.39 mmol) in THF (0.70 ml)  
25 was cooled to 0°C under N<sub>2</sub>. A solution of Normant reagent (~ 0.5 M) in THF was added in 1 ml portions until the reaction was complete by TLC (10/90 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>). The reaction was quenched with NH<sub>4</sub>Cl (sat), diluted with CH<sub>2</sub>Cl<sub>2</sub>, and 1 N NaOH was added. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the combined organics were dried over  
30 K<sub>2</sub>CO<sub>3</sub>, filtered, and concentrated to yield **150** (0.17 g, 71% yield). LRMS calc'd: 602; found (M+H): 603.

#### Example 47

**485**

-108-



4

**Step 1:** Ethyl 3-hydroxy-4-iodobenzoate (2.92 g), sodium 3,4-methylenedioxybenzenesulfinate (3.2 g), copper (I) iodide (2.8 g) and DMF (20 ml) were heated under  $\text{N}_2$  for 20 h, added to 20% aqueous  $\text{NaI}$  (200 ml), extracted with  $\text{EtOAc}$ , dried and evaporated. The residue was flash-chromatographed on silica, eluting with hexanes- $\text{EtOAc}$ , and pure fractions evaporated to give the product (120) as a solid (0.81 g), mp 133-135°C.

**Step 2** (120) (0.40 g) in THF (5 ml) was added to a stirred, ice-cooled mixture of  $\text{LiAlH}_4$  (0.08 g) and THF (10 ml). After 0.5 h, the mixture was treated with  $\text{H}_2\text{O}$  then 1N  $\text{HCl}$ , extracted with  $\text{EtOAc}$ , dried over  $\text{MgSO}_4$  and evaporated to give (121), which was used in the next step.

**Step 3:** (121) in  $\text{CH}_2\text{Cl}_2$  (20 ml) and  $\text{Et}_3\text{N}$  (1.0 ml) was stirred with ice cooling and  $\text{CH}_3\text{SO}_2\text{Cl}$  (0.21 ml) added. After 0.5 h, the mixture was washed with 1N  $\text{HCl}$ , and then  $\text{NaHCO}_3$  solution, dried and evaporated. This residue was stirred with diisopropylethylamine (0.3 ml) and the piperazine derivative (0.5 g) in DMF (3 ml) for 20 h at RT, worked up in water- $\text{EtOAc}$ , dried, evaporated and




-109-

chromatographed on flash silica, eluting with 1 to 3% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>. Pure fractions were evaporated to give the product (122) as a white foam (0.60 g). Mass spectrum: MH<sup>+</sup> = 638.


5 Step 4: (122) (0.43 g) was stirred for 45 min at RT in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) and 95% TFA (5 ml), evaporated, worked up in EtOAc - aq, NaHCO<sub>3</sub>, dried, and evaporated to give (123) as a white foam (0.27 g). Mass spectrum: MH<sup>+</sup> = 538.

10 Step 5: (123) (0.07 g) and o-toluoyl chloride (0.06 g) were stirred for 45 min at RT in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) and saturated NaHCO<sub>3</sub> (5 ml). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the organics were evaporated. The residue was stirred for 0.5 h in a 5% solution of NaOH in 95% CH<sub>3</sub>OH (4 ml), diluted with H<sub>2</sub>O and CH<sub>2</sub>Cl<sub>2</sub>, and treated with stirring with small portions of solid CO<sub>2</sub> until the pH of the aq. phase was 7-8. The organic phase was dried and evaporated, and the residue dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) and added to Et<sub>2</sub>O (15 ml) containing 4M HCl-dioxan (0.4 ml). The precipitate was centrifuged, the solid washed twice with Et<sub>2</sub>O, and dried at RT in N<sub>2</sub> to give the hydrochloride of 485 as a white powder (0.051 g). Mass spectrum: MH<sup>+</sup> = 607.

20

**Example 48**

-110-



**Step 1:** The Grignard reagent prepared by reacting 3,4-methylenedioxy bromobenzene (20 g) and Mg (3 g) in THF (60 ml) was cooled in ice and 3-benzyloxybenzaldehyde (16 g) in THF (60 ml) was added slowly. After 15 min, the mixture was added to a stirred mixture of ice and HCl (75 ml of 2N), extracted with Et<sub>2</sub>O, washed with NaHCO<sub>3</sub> solution, dried and evaporated. The residue was extracted by stirring followed by decanting with hexanes (2 x 100 ml) and the residue was dried at high vacuum to a brown oil.

This material was shaken in H<sub>2</sub> (60 psi) at RT for 4 h in EtOAc (150 ml), HOAc (1.5 ml) and 20% palladium hydroxide on carbon (1.5 g). After filtration and evaporation, the residue was triturated with 1:1 Et<sub>2</sub>O-hexanes (2 x 50 ml) and dried at RT to give (124) as a white solid (8.2 g). Mp 139-142°C.

**Step 2:** (124) (8.1 g) was stirred with Et<sub>3</sub>SiH (20 ml) in CH<sub>2</sub>Cl<sub>2</sub> (200 ml) and TFA (10 ml) was added dropwise. After 0.5 h. the solution was washed twice with H<sub>2</sub>O and evaporated. The residue in CH<sub>2</sub>Cl<sub>2</sub> (100 ml) was treated for 15 min. with 1M tetrabutylammonium fluoride in THF (50 ml), washed with 1N H<sub>2</sub>SO<sub>4</sub>, dried, evaporated and chromatographed on flash silica, eluting with 0 to 2% Et<sub>2</sub>O in CH<sub>2</sub>Cl<sub>2</sub>. Pure fractions on evaporation gave (125) as a thick oil (6.28 g). Mass spectrum: MH<sup>+</sup> = 229.


**Step 3:** A mixture of (125) (2.3 g), 4-t-butoxycarbonyl-2-(R)-methylpiperazine (2.4 g), ethanol (40 ml), 37% formalin (2.5 ml) and 4M HCl-dioxan (1.25 ml) was refluxed for 24 h, worked up in CH<sub>2</sub>Cl<sub>2</sub>: aq. NaHCO<sub>3</sub>, dried, evaporated and the residue was chromatographed on flash silica, eluting with 5 to 30% Et<sub>2</sub>O-hexanes. Pure fractions were evaporated to give (126) as a white foam (2.6 g). Mass spectrum: MH<sup>+</sup> = 441.

**Step 4:** A solution of (126) (1.7 g) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml), H<sub>2</sub>O (0.5 ml) and TFA (5 ml) was stirred at RT for 1 h., diluted with CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O, and treated with small portions of K<sub>2</sub>CO<sub>3</sub> until the pH of the

-111-

aqueous phase remained at 8-9. The organic phase was dried and evaporated to give (127) as a white foam (1.2 g). Mass spectrum:  $MH^+ = 341$ .

**Step 5:** A mixture of (127) (1.12 g), 1-(o-toluoyl)-4-piperidinone (0.8 g),  $CH_2Cl_2$  (25 ml) and  $NaBH(OAc)_3$  (1.5 g) was stirred at RT for 20 h, washed with excess aq.  $NaHCO_3$ , dried and evaporated. The residue was dissolved in  $CH_3OH$  containing 4M HCl-dioxan (2 ml), evaporated, co-evaporated with  $CH_3OH$  (100 ml), and this residue dissolved in  $CH_3OH$  (1 ml) and  $CH_2Cl_2$  (15 ml) and added to stirred  $Et_2O$  (100 ml) containing 4M HCl-dioxan (1 ml). The precipitate was filtered, washed with  $Et_2O$  and dried to give the hydrochloride of **300** as a white powder (1.76 g). Mass spectrum:  $MH^+ = 542$ .

**Example 49**

**Step 1:** A mixture of 3-iodophenol (1.76 g), the piperazine (1.35 g), 37% formalin (1.25 ml) and ethanol (15 ml) was refluxed for 1 h, then additional formalin (0.75 ml) was added and heating continued for

-112-

10h. The reaction was cooled, worked up in  $\text{CH}_2\text{Cl}_2$ - $\text{H}_2\text{O}$ , dried and evaporated. The major product was isolated by flash chromatography on silica, eluting with 0 to 1.5%  $\text{CH}_3\text{OH}$  in EtOAc. Evaporation of pure fractions gave a yellow foam, (128) (1.31 g).

5 Mass spectrum:  $\text{MH}^+ = 502$ .

Step 2: (128) (1.25 g) and t-butylchlorodimethylsilane (0.8 g) with  $\text{Et}_3\text{N}$  (0.8 ml) and dimethylaminopyridine (0.03 g) was stirred in  $\text{CH}_2\text{Cl}_2$  (30 ml) for 40h, evaporated, and chromatographed on flash silica in EtOAc to obtain the product (129) as a white foam (1.48 g).

10 Mass spectrum:  $\text{MH}^+ = 602$ .

Step 3: A solution of (129) (1.42 g) in dry THF (25 ml) at  $-70^\circ\text{C}$  was treated with n-BuLi in hexanes (2M; 1.4 ml). Immediately, a solution of 3,4-methylenedioxybenzenesulfonyl fluoride (0.65 g) in THF (1 ml) was added, and the mixture stirred to RT over 0.5h. The reaction was partitioned in  $\text{H}_2\text{O}$  - EtOAc, dried and evaporated, and the residue chromatographed on flash silica in hexanes-EtOAc. Pure fractions were evaporated to give the product (130) as a pale yellow foam (0.76 g). Mass spectrum:  $\text{MH}^+ = 674$ .

15 Step 4: (130) (0.55 g) was stirred for 20 h in  $\text{CH}_3\text{OH}$  (15 ml) containing HOAc (0.1 g) and KF (1.0 g), partitioned in  $\text{H}_2\text{O}$ - $\text{CH}_2\text{Cl}_2$ , dried and evaporated. Flash chromatography on silica with EtOAc gave the pure product (131) as a pale yellow foam (0.41 g). Mass spectrum:  $\text{MH}^+ = 560$ .

20 Step 5: A mixture of (131) (0.37 g),  $\text{CH}_2\text{Cl}_2$  (3 ml),  $\text{H}_2\text{O}$  (0.05 ml) and TFA (2 ml) was kept at RT for 2 h, evaporated, and partitioned in  $\text{H}_2\text{O}$ - $\text{CH}_2\text{Cl}_2$  with sufficient  $\text{NaHCO}_3$  to maintain the aq. phase at pH 8-9. The  $\text{CH}_2\text{Cl}_2$  extracts were dried and evaporated, and this residue in DMF (15 ml) stirred 20 h at RT with o-toluic acid (0.2 g), 1-hydroxybenzotriazole (0.15 g) and EDCI (0.4 g). The mixture was partitioned in EtOAc and aq.  $\text{NaHCO}_3$ , dried and evaporated. The major component was isolated by ptlc on silica plates with 3%  $\text{CH}_3\text{OH}$ - $\text{CH}_2\text{Cl}_2$  to give a white foam (0.38 g) which was mainly the N,O-di-toluoyl compound. 0.3 g of this material was stirred for 24 h at RT in  $\text{CH}_3\text{OH}$  (7 ml) and  $\text{H}_2\text{O}$  (0.8 ml) with NaOH (0.5 g). The solution was stirred with  $\text{H}_2\text{O}$ - $\text{CH}_2\text{Cl}_2$  and neutralised to pH 7-8 with

25


30

35

-113-


solid CO<sub>2</sub>. The organic phase was dried and evaporated, and the hydrochloride of **210** precipitated as described in earlier examples, and dried at RT in vacuum to give a white solid (0.22 g). Mass spectrum: MH<sup>+</sup> = 578.

5

**Example 50**

10

27 mg of **30** were taken up in 1.5 mL of glacial HOAc. The solution was stirred for 7 h at 40°C. Excess saturated NaHCO<sub>3</sub> solution was added, and the mixture was extracted with EtOAc. This was dried over Na<sub>2</sub>SO<sub>4</sub>, evaporated, and the residue purified by column chromatography in 10% CH<sub>3</sub>OH/EtOAc, to give approx. 17 mg of **520** as an oil. MS (FAB): 598.4 (M+1), 538.5.

**Example 51**

15


-114-

To a solution of (132) (2.31 g, 6.00 mmol) in anhydrous  $\text{CH}_3\text{CN}$  (30 ml) were successively added 4-piperidone ethylene ketal (0.77 ml, 6.00 mmol) and 2,2,6,6-tetramethyl piperidine (1.22 ml, 7.20 mmol), and the mixture was heated 14 h at  $60^\circ\text{C}$ . After evaporation of the solvent, the residue was taken up in saturated aqueous  $\text{NaHCO}_3$  (100 ml), extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 50 ml), dried over  $\text{Na}_2\text{SO}_4$ , and evaporated. The oil thus obtained (2.89 g) was refluxed with 6 N  $\text{HCl}$  (40 ml) for 7 h then evaporated. Workup as above and purification by flash chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_3$  96:4:1) afforded (133) (1.26 g, 49%).

































To a solution of (133) (470 mg, 1.21 mmol) and *N*-Boc piperazine (255 mg, 1.27 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (1 ml) was added titanium (IV) isopropoxide (447 ml, 1.21 mmol), and the mixture was stirred 3 days at room temperature. The solution was diluted with  $\text{CH}_2\text{Cl}_2$  (1 ml),  $\text{Et}_2\text{AlCl}$  1 N in toluene (3.65 ml) was added at  $0^\circ\text{C}$  and the resulting mixture was allowed to warm to room temperature and stirred 2 h. Upon addition of Celite and water (100 ml), the final suspension was filtered over Celite, dried over  $\text{Na}_2\text{SO}_4$  and evaporated. The crude thus obtained (655 mg, 93%) was dissolved in anhydrous THF (7 ml), treated at  $0^\circ\text{C}$  with  $\text{CH}_3\text{MgBr}$  3 N in  $\text{Et}_2\text{O}$  (3.8 ml, 11.3 mmol) and the solution was allowed to warm to room temperature then heated 3 h at  $55^\circ\text{C}$ . After cooling, Celite (2 g) and  $\text{CH}_2\text{Cl}_2$  (10 ml) were added and the suspension was slowly poured into ice. The resulting suspension was filtered over Celite and worked up as before to provide (134) (375 mg, 58%), after flash chromatography ( $\text{SiO}_2$ , eluent  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_3$  80:20:1).

Removal of the Boc group with TFA gave, after basic workup with 1 N  $\text{NaOH}$  and extraction with  $\text{CH}_2\text{Cl}_2$ , 260 mg of intermediate. This (20 mg, 0.042 mmol) was subjected to *N*-acylation with *o*-toluoyl chloride to provide, after conversion to the hydrochlorid salt, 587 (15 mg) as a white solid:  $\text{MH}^+ = 590.2$ .































Using the appropriate starting materials in the procedures described above or modifications of those procedures well known to those skilled in the art, the compounds shown in the following tables are prepared.

































No.	R	W	Physical Data
151			m.p. = 198-206°C
152			m.p. = 186-192°C
153			(M-15) = 596
154			m.p. = 210-15°C
155			MH+ = 612
156			MH+ = 596
157			MH+ = 660
158			MH+ = 626
159			MH+ = 610
160			MH+ = 650
161			HRMS calc'd: 614.2664 found: 614.2664
162			HRMS calc'd: 560.2947 found: 560.2955
163			Mass spec. (FAB)M+1=635
164			Mass spec. (FAB)M+1=687
165			Mass spec. (FAB)M+1=656

166			Mass spec. (FAB)M+1=580
167			HRMS calc'd: 560.2947 found: 560.2933
168			HRMS calc'd: 610.1738 found: 610.1739
169			HRMS calc'd: 546.2790 found: 546.2787
170			HRMS calc'd: 588.2896 found: 588.2897
171			HRMS calc'd: 624.2896 found: 624.2903
172			HRMS calc'd: 628.1645 found: 628.1646
173			HRMS calc'd: 600.2696 found: 600.2694
174			HRMS calc'd: 541.2285 found: 541.2289
175			HRMS calc'd: 646.1551 found: 646.1150
176			HRMS calc'd: 559.2191 found: 559.2202
177			HRMS calc'd: 604.2845 found: 604.2840
178			HRMS calc'd: 640.2845 found: 640.2840
179			HRMS calc'd: 668.1794 found: 668.1785
180			HRMS calc'd: 580.2401 found: 580.2401
181			HRMS calc'd: 616.2401 found: 616.2395




182			FAB(MH+) = 606
183			FAB(MH+) = 648
184			FAB(MH+) = 632
185			FAB(MH+) = 606
186			FAB(MH+) = 652
187			FAB(MH+) = 636
188			FAB(MH+) = 636
189			FAB(MH+) = 686
190			m.p. = 200-202°C (dec)
191			m.p. = 201-203°C (dec)
192			m.p. = 215-216°C (dec)
193			m.p. = 210-201°C (dec)
194			m.p. = 219-220°C (dec)
195			FAB(MH+) = 582
196			m.p. = 204-206°C (dec)































197			m.p. = 198-200°C (dec)
198			MH+ = 596
199			MH+ = 610
200			MH+ = 620
201			MH+ = 606
202			MH+ = 541
203			MH+ = 591
204			MH+ = 620
205			MH+ = 620
206			MH+ = 620
207			MH+ = 605
208			MH+ = 592
209			MH+ = 520
211			MH+ = 684
212			MH+ = 621

















































-119-































213			MH+ = 634
214			m.p.= decomp >225°C
215			m.p.= decomp 184°C
216			m.p.= decomp above 189°C
217			m.p.= 172-176°C































































No.	R	W	R <sup>1</sup>	R <sup>27</sup>	X	Physical Data
218			H	H	-SO-	MS (electrospray): 546 (M+1), 487 (isomer A)
219			H	H	-SO-	MS (electrospray): 546 (M+1), 463 (isomer B)
220			H	H	-SO <sub>2</sub> -	MS (FAB): 562 (M+1), 302
221			H	H	-SO <sub>2</sub> -	MS (FAB): 548 (M+1), 461
222			H	H	-SO <sub>2</sub> -	MS (electrospray): 548.2 (M+1), 465
223			H	H	-SO <sub>2</sub> -	MS (electrospray): 534.2 (M+1), 472
224			H	H	-SO <sub>2</sub> -	MS (electrospray): 613, 615 (M+1), 472





























225			H	H	-SO <sub>2</sub> -	m.p.= 230-246°C
226			H	H	-SO <sub>2</sub> -	m.p.= 220°C (dec)
227			H	H	-SO <sub>2</sub> -	m.p.= decomp above 245°C
228			H	H	-SO <sub>2</sub> -	m.p.= 169-171°C
229			H	H	-SO <sub>2</sub> -	m.p.= decomp above 235°C
230			H	H	-SO <sub>2</sub> -	m.p.= decomp above 202°C
231			H	H	-SO <sub>2</sub> -	m.p.= decomp above 208°C
232			H	H	-SO <sub>2</sub> -	m.p.= decomp above 195°C
233			H	H	-SO <sub>2</sub> -	m.p.= decomp above 235°C
234			H	H	-SO <sub>2</sub> -	m.p.= decomp above 217°C
235			H	H	-SO <sub>2</sub> -	m.p.= decomp above 185°C
236			H	H	-SO <sub>2</sub> -	m.p.= decomp above 207°C
237			H	H	-SO <sub>2</sub> -	m.p.= 164-166°C
238			H	H	-SO <sub>2</sub> -	m.p.= decomp above 250°C
239			H	H	-SO <sub>2</sub> -	m.p.= decomp above 239°C

240			H	H	-SO <sub>2</sub> -	m.p.= 204 <sup>o</sup> C
241			H	H	-SO <sub>2</sub> -	m.p.= decomp above 232 <sup>o</sup> C
242			H	H	-SO <sub>2</sub> -	m.p.= decomp above 205 <sup>o</sup> C
243			H	H	-SO <sub>2</sub> -	m.p.= decomp above 180 <sup>o</sup> C
244			H	H	-SO <sub>2</sub> -	m.p.= decomp above 200 <sup>o</sup> C
245				H	-SO <sub>2</sub> -	MH <sup>+</sup> = 576
246				H	-SO <sub>2</sub> -	MH <sup>+</sup> = 596
247			H		-SO <sub>2</sub> -	m.p.= 174-177 <sup>o</sup> C
248				H		MH <sup>+</sup> = 542
249			H	H		FAB(MH <sup>+</sup> ) = 526
250						HRMS calc'd: 544.2731 found: 544.2724
251						Mass. Spec. (FAB)M <sup>+</sup> 1 = 548
252						Mass. Spec. (FAB)M <sup>+</sup> 1 = 601
253				H	-CH <sub>2</sub> -	MH <sup>+</sup> = 526
254					-CH <sub>2</sub> -	HRMS calc'd: 554.3383 found: 554.3366

255			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 530.2938 found: 530.2943
256			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 556.3539 found: 556.3547
257			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	Mass. Spec. (FAB)M+1 = 538
258			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 554.3383 found: 554.3366
259			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 552.3590 found: 552.3602
260			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 544.2998 found: 544.2987
261			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 524.3641 found: 524.3647
262			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 510.3484 found: 510.3476
263			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 574.2433 found: 574.2414
264			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 511.3437 found: 511.3448
265			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 527.3386 found: 527.3386
266			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 526.3434 found: 526.3442
267			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 590.2382 found: 590.2391
268			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 540.3590 found: 540.3611
269			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	HRMS calc'd: 502.2892 found: 502.2898
























-123-











270					-CH <sub>2</sub> -	HRMS calc'd: 566.1841 found: 566.1839
271					-CH <sub>2</sub> -	HRMS calc'd: 479.2481 found: 479.2477
272					-CH <sub>2</sub> -	HRMS calc'd: 530.2938 found: 530.2954
273					-CH <sub>2</sub> -	HRMS calc'd: 574.3434 found: 574.3436
274					-CH <sub>2</sub> -	HRMS calc'd: 514.3234 found: 514.3233
275					-CH <sub>2</sub> -	HRMS calc'd: 550.3234 found: 550.3218
276					-CH <sub>2</sub> -	HRMS calc'd: 502.2892 found: 502.2893
277					-CH <sub>2</sub> -	HRMS calc'd: 479.2481 found: 479.2469
278					-CH <sub>2</sub> -	HRMS calc'd: 538.2892 found: 538.2884
279					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 542
280					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 556
281					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 528
282					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 556
283					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 602
284					-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 578


285			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 579
286			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 591
287			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	FAB(MH <sup>+</sup> ) = 611
288			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	m.p. = 215-216°C (dec)
289			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	m.p. = 178-179°C (dec)
290			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	m.p. = 209-211°C (dec)
291			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 530
292			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 541
293			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 546
294			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 561
295			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 556
296			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 527
297			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 527
298			$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \text{---} \\ \text{---} \end{array}$	-CH <sub>2</sub> -	MH <sup>+</sup> = 526

















299			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 541$
301			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 541$
303			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 570$
304			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 555$
305			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 543$
306			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{MH}^+ = 542$
307			H	H	$-\text{CH}_2-$	m.p. = 216-223°C
308			H	$\text{CH}_3$ $\text{CH}_3$	$-\text{CH}_2-$	m.p. = decomp. above 145°C
309			H	$\text{CH}_3$ ▼	$-\text{CH}_2-$	m.p. = decomp. above 110°C
309 A			H	$\text{CH}_3$	$-\text{CH}_2-$	m.p. = 228°C (dec)
494			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{FAB}(\text{MH}^+) = 546$
495			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	$\text{FAB}(\text{MH}^+) = 576$
496			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	m.p. = 195-201 °C
497			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	m.p. = 202- 204°C

498			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	m.p. = 202-204°C
499			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	m.p. = 228-232°C
500			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	MH+ = 562
501			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_2-$	MH+ = 560
521			H	$\text{CH}_3$	$-\text{O}-$	m.p. = 197-199°C
522			H	$\text{CH}_3$	$-\text{O}-$	m.p. = 175-177°C
523			H	$\text{CH}_3$	$-\text{O}-$	m.p. = 195-197°C
524			H	$\text{CH}_3$	$-\text{O}-$	m.p. = 60-62°C
525			H	$\text{CH}_3$	$-\text{O}-$	m.p. = 196-200°C (isomer A)
526			H	$\text{CH}_3$	$-\text{O}-$	FAB(MH+) = 523 (isomer B)
527	$\text{H}_3\text{CO}-$		H	H	$\text{O}-\text{C}-$	HRMS (FAB) calc'd (MH+): 450.2757 Found: 450.2752 m.p. = 245-260°C
559			H	H	$-\text{SO}_2-$	m.p. = decomp above 115°C


560			$\text{CH}_3$	$\text{CH}_3$	$\begin{array}{c} \text{OH} \\   \\ -\text{C}- \\   \\ \text{H} \end{array}$	m.p. = 161-166°C
561			$\text{CH}_3$	$\text{CH}_3$	$\begin{array}{c} \text{OH} \\   \\ -\text{C}- \\   \\ \text{H} \end{array}$	Mass. Spec. (FAB)M+1 = 635
562			$\text{CH}_3$	$\text{CH}_3$	-CH <sub>2</sub> -	HRMS calc'd: 618.2331 found: 618.2316
592			-CF <sub>3</sub>	H	-SO <sub>2</sub> -	HRMS calc'd: 664.1456 found: 664.1452
593			$\text{CH}_3$	$\text{CH}_3$	$\begin{array}{c} \text{OH} \\   \\ -\text{C}- \\   \\ \text{H} \end{array}$	HRMS calc'd: 566.3183 found: 566.3180



No.	R	W	R <sup>1</sup>	R <sup>27</sup>	R <sup>20</sup>	Physical Data
310			$\text{CH}_3$	$\text{CH}_3$	-CH <sub>3</sub>	MH <sup>+</sup> = 603
311			$\text{CH}_3$	$\text{CH}_3$	-CH <sub>3</sub>	MH <sup>+</sup> = 599
312			$\text{CH}_3$	$\text{CH}_3$	-CH <sub>3</sub>	MH <sup>+</sup> = 597
313			$\text{CH}_3$	$\text{CH}_3$		MH <sup>+</sup> = 626
314			$\text{CH}_3$	$\text{CH}_3$		MH <sup>+</sup> = 655
315			$\text{CH}_3$	$\text{CH}_3$	n-Pr	MH <sup>+</sup> = 611



















































316			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	MH <sup>+</sup> = 584
317			H	H	$-\text{CH}_3$	HRMS calc'd: 555.2971 found: 555.2983 m.p.=235-240°C
318			H	H	$-\text{CH}_3$	HRMS calc'd: 569.3128 found: 569.3132 m.p.=170-175°C
319			H	H	$-\text{CH}_3$	HRMS calc'd: 559.2721 found: 559.2732 m.p.=192-198°C
320			H	H	$-\text{CH}_3$	HRMS calc'd: 575.2425 found: 575.2418 m.p.=205-210°C
321			H	H	$-\text{CH}_3$	HRMS calc'd: 619.20 found: 619.1911 m.p.=170-180°C
322			H	H	$-\text{CH}_3$	HRMS calc'd: 555.3355 found: 555.3355 m.p.=175-185°C
323			H	H	$-\text{CH}_3$	HRMS calc'd: 555.3335 found: 555.3330 m.p.=180-190°C
324			H	H	$-\text{CH}_3$	HRMS calc'd: 525.3230 found: 525.3224 m.p.=205-210°C
325			H	H	$-\text{CH}_3$	HRMS calc'd: 539.3386 found: 539.3397 m.p.=228-237°C
326			H	H	$-\text{CH}_3$	HRMS calc'd: 539.3386 found: 539.3396 m.p.=234-239°C











































327			H	H	-CH <sub>3</sub>	HRMS calc'd: 591.2971 found: 591.2966 m.p.=195-200°C
329			H	H	-CH <sub>3</sub>	m.p. - 185-190°C (dec)
330			H	H	-CH <sub>3</sub>	m.p. - 240-248°C (dec)
331			H	H	-CH <sub>3</sub>	m.p. - 235-240°C (dec)
332			H	H	-CH <sub>3</sub>	m.p. - 185-195°C (dec)
333			H	H	-CH <sub>3</sub>	m.p. - 175-195°C (dec)
334			H	H	-CH <sub>3</sub>	m.p. - 230-235°C (dec)
335			H	H	-CH <sub>3</sub>	m.p. - 245-255°C (dec)
336			H	H	-CH <sub>3</sub>	m.p. - 250-255°C (dec)










































No.	R	W	R <sup>1</sup> , R <sup>27</sup>	X	Physical Data
337				-CH <sub>2</sub> -	LRMS: calc'd: 582 found: 583
338				-CH <sub>2</sub> -	LRMS: calc'd: 538 found: 539
339				-CH <sub>2</sub> -	LRMS: calc'd: 574 found: 575



























340				-CH <sub>2</sub> -	LRMS: calc'd: 584 found: 585
341			=O	-CH <sub>2</sub> -	LRMS: calc'd: 540 found: 541
342			H, OH	-CH <sub>2</sub> -	LRMS: calc'd: 526 found: 527
436				-SO <sub>2</sub> -	LRMS: calc'd: 632 found: 633
437				-SO <sub>2</sub> -	LRMS: calc'd: 646 found: 647
438				-SO <sub>2</sub> -	LRMS: calc'd: 558 found (M+H): 559
439				-SO <sub>2</sub> -	LRMS: calc'd: 572 found (M+H): 573
440				-SO <sub>2</sub> -	LRMS (FAB): calc'd (M+H): 591 found: 591
441				-SO <sub>2</sub> -	LRMS (FAB): calc'd: 600 found (M+H): 601
442				-SO <sub>2</sub> -	LRMS (FAB): calc'd: 626 found (M+H): 627
443				-CH <sub>2</sub> -	LRMS: calc'd: 592 found (M+H): 593
444				-SO <sub>2</sub> -	LRMS: calc'd: 626 found (M+H): 627
445				-SO <sub>2</sub> -	LRMS: calc'd: 634 found (M+H): 635
446				-SO <sub>2</sub> -	LRMS: calc'd: 620 found (M+H): 621
447				-SO <sub>2</sub> -	LRMS: calc'd: 646 found (M+H): 647
448				-SO <sub>2</sub> -	LRMS: calc'd: 606 found (M+H): 607
449				-SO <sub>2</sub> -	LRMS: calc'd: 660 found (M+H): 661

450				-SO <sub>2</sub> -	LRMS: calc'd: 652 found (M+H): 653
451				-SO-	LRMS: calc'd: 592 found (M+H): 593
452				-SO-	LRMS: calc'd: 622 found (M+H): 623
453				-SO-	LRMS: calc'd: 618 found (M+H): 619
454				-SO-	LRMS: calc'd: 602 found (M+H): 603
455				-SO-	LRMS: calc'd: 630 found (M+H): 631
456				-SO <sub>2</sub> -	LRMS: calc'd: 650 found (M+H): 651
457				-SO <sub>2</sub> -	LRMS: calc'd: 638 found (M+H): 639
458				-SO-	LRMS: calc'd: 634 found (M+H): 635
459				-S-	LRMS: calc'd: 586 found (M+H): 587
460				-S-	LRMS: calc'd: 574 found (M+H): 575
461				-S-	LRMS: calc'd: 614 found (M+H): 615
462				-S-	LRMS: calc'd: 618 found (M+H): 619
463				-S-	LRMS: calc'd: 606 found (M+H): 607
464				-S-	LRMS: calc'd: 602 found (M+H): 603
465				-SO <sub>2</sub> -	LRMS: calc'd: 596 found (M+H): 597
466		H		-SO <sub>2</sub> -	LRMS: calc'd: 500 found (M+H): 501
















467				-SO <sub>2</sub> -	LRMS: calc'd: 691 found (M+H): 692
468				-SO <sub>2</sub> -	LRMS: calc'd: 705 found (M+H): 706
469				-SO <sub>2</sub> -	LRMS: calc'd: 662 found (M+H): 663
470				-SO <sub>2</sub> -	LRMS: calc'd: 620 found (M+H): 621
471				-SO <sub>2</sub> -	LRMS: calc'd: 654 found (M+H): 655
472				-S-	LRMS: calc'd: 584 found (M+H): 585
473				-SO-	LRMS: calc'd: 600 found (M+H): 601
474				-SO <sub>2</sub> -	LRMS: calc'd: 616 found (M+H): 617
475				-SO-	LRMS: calc'd: 618 found (M+H): 619
476				-SO <sub>2</sub> -	LRMS: calc'd: 634 found (M+H): 635
477				-SO <sub>2</sub> -	LRMS: calc'd: 592 found (M+H): 593
478				-SO <sub>2</sub> -	LRMS: calc'd: 620 found (M+H): 621
479				-SO <sub>2</sub> -	LRMS: calc'd: 604 found (M+H): 605
480				-SO <sub>2</sub> -	LRMS: calc'd: 632 found (M+H): 633



481				-SO <sub>2</sub> -	LRMS: calc'd: 625 found (M+H): 625
482				-SO <sub>2</sub> -	LRMS: calc'd: 636 found (M+H): 637
483				-SO-	LRMS: calc'd: 576 found (M+H): 577
484				-SO-	LRMS: calc'd: 588 found (M+H): 589
488			=O	-SO <sub>2</sub> -	LRMS: calc's: 590 found: 591
489			=O	-SO <sub>2</sub> -	LRMS: calc's: 580 found: 581
490			H, OH	-SO <sub>2</sub> -	LRMS: calc's: 576 found: 577
491			CH <sub>3</sub> , OH	-SO <sub>2</sub> -	LRMS: calc's: 590 found: 591
492				-SO <sub>2</sub> -	LRMS: calc's: 618 found: 619
493			=O	-S-	LRMS: calc's: 542 found (M+H): 543
502			H, H	-SO <sub>2</sub> -	LRMS: calc's: 566 found: 567
503			H, H	-SO <sub>2</sub> -	LRMS: calc's: 576 found: 577
504			H, H	-SO <sub>2</sub> -	LRMS: calc's: 546 found: 547
505			H, H	-SO <sub>2</sub> -	LRMS: calc's: 581 found: 581
506			H, H	-SO <sub>2</sub> -	LRMS: calc's: 625 found: 626
507			H, H	-SO <sub>2</sub> -	LRMS: calc's: 564 found: 565
508			H, H	-SO <sub>2</sub> -	LRMS: calc's: 630 found: 631


509			H, H	-SO <sub>2</sub> -	LRMS: calc's: 546 found: 547
510			H, H	-SO <sub>2</sub> -	LRMS: calc's: 560 found: 561
511			H, H	-SO <sub>2</sub> -	LRMS: calc's: 552 found: 553
512			H, H	-SO <sub>2</sub> -	LRMS: calc's: 567 found: 567
513			H, H	-SO <sub>2</sub> -	LRMS: calc's: 611.6 found: 613
514			H, H	-SO <sub>2</sub> -	LRMS: calc's: 562 found: 563
515			H, H	-SO <sub>2</sub> -	LRMS: calc's: 672 found: 673
516			H, H	-SO <sub>2</sub> -	LRMS: calc's: 582 found: 583
517			H, H	-SO <sub>2</sub> -	LRMS: calc's: 596 found: 597
518			H, H	-SO <sub>2</sub> -	LRMS: calc's: 624 found: 625
519			H, H	-SO <sub>2</sub> -	LRMS: calc's: 596 found (M+H <sup>+</sup> ): 597
563	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-S-	HRMS: calc'd (M+1): 488.2736 found: 488.2728
564	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-SO <sub>2</sub> -	HRMS: calc'd (M+1): 519.2681 found: 519.2685
565	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-SO <sub>2</sub> -	HRMS: calc'd (M+1): 483.2681 found: 483.2688
566	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-SO <sub>2</sub> -	HRMS: calc'd (M+1): 503.2135 found: 503.2144

-135-











567	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-S-	HRMS: calc'd (M+1): 451.2873 found: 451.2870
568	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-CH <sub>2</sub> -	HRMS: calc'd (M+1): 469.3219 found: 469.3219
569	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-S-	HRMS: calc'd (M+1): 489.2688 found: 489.2687
570	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-S-	HRMS: calc'd (M+1): 488.2736 found: 488.2739
571			H, H	-O-	HRMS: calc'd (M+1): 424.2600 found: 424.2608
572	-CH <sub>3</sub>		H, H	-O-	HRMS: calc'd (M+1): 384.2287 found: 384.2294
573	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-S-	HRMS: calc'd (M+1): 488.2736 found: 488.2751
574	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-O-	HRMS: calc'd (M+1): 472.2964 found: 472.2964
575	(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -		H, H	-SO-	HRMS: calc'd (M+1): 503.2732 found: 503.2723
576	CH <sub>3</sub> CH <sub>2</sub> SO <sub>2</sub> -		H, H	-NH-	LRMS: calc'd: 519 found (M+H <sup>+</sup> ): 520
577	-CH <sub>3</sub>		H, H	-NH-	LRMS: calc'd: 441 found (M+H <sup>+</sup> ): 442
578				=O	LRMS: calc'd: 564 found: 565

-136-


579					LRMS: calc'd: 754 found: 755
580				=O	LRMS: calc'd: 603 found: 603
581		-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>		=O	LRMS: calc'd: 570 found: 571
582				=O	LRMS: calc'd: 596 found: 597
583				=O	LRMS: calc'd: 610 found: 611
584					LRMS: calc'd: 579 found: 580
585			H, OH	=O	LRMS: calc'd: 528 found: 529
586			H, H	=O	LRMS: calc'd: 512 found: 513
590			H, H	-SO <sub>2</sub> -	LRMS: calc'd: 574 found: 575
591			H, H	-O-	HRMS: calc'd (M+1): 483.3012 found: 483.3008























































No.	X	R <sup>1</sup>	R <sup>27</sup>	Physical Data
533		H	H	MS (IFAB): 555.4 (M+1), 391,307

534		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 569.4 (M+1), 553, 302
535		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 583.4 (M+1), 302, 282
536		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 611 (M+1), 554 (isomer A)
537		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 611 (M+1), 325 (isomer B)
538		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 626.4 (M+1), 554
539		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 610.3 (M+1), 460 (isomer A)
540		$\text{CH}_3$	$\text{CH}_3$	MS (fFAB): 610.2 (M+1), 554, 309 (isomer B)
541		$\text{CH}_3$	$\text{CH}_3$	MS (electrospray): 620 (M+1), 551, 433 (isomer A)
542		$\text{CH}_3$	$\text{CH}_3$	MS (electrospray): 620 (M+1), 551, 433 (isomer B)
543		H	H	MS (fFAB): 599.4 (M+1), 510, 277

























































544				MS (fFAB): 627.4 (M+1), 326, 302
545				MS (fFAB): 597.5 (M+1), 302
546		H	H	MS (electrospray): 555 (M+1), 510, 392
547		H	H	MS (fFAB): 581.4 (M+1), 510
548		H	H	MS (fFAB): 667.4 (M+1), 510
549		H	H	MS (electrospray): 619.2 (M+1), 510
550		H	H	FAB(MH+) = 611
551		H	H	MS (electrospray): 624.2 (M+1), 510
552		H	H	MS (electrospray): 599.2 (M+1), 510













































































No.	R	W	R <sup>1</sup>	R <sup>27</sup>	R <sup>30</sup>	Physical Data
343			H	H	-CH <sub>3</sub>	MS (electrospray): 524.1 (M+), 373
344					-CH <sub>3</sub>	MH <sup>+</sup> = 579
345					-CH <sub>3</sub>	MH <sup>+</sup> = 593
346					-CH <sub>3</sub>	MH <sup>+</sup> = 641
347					-CH <sub>3</sub>	MH <sup>+</sup> = 655
348					-CH <sub>3</sub>	MH <sup>+</sup> = 669
349					-CH <sub>3</sub>	FAB (MH <sup>+</sup> ) = 618
350					-CH <sub>3</sub>	FAB (MH <sup>+</sup> ) = 610
351					-CH <sub>3</sub>	FAB (MH <sup>+</sup> ) = 620
352					-CH <sub>3</sub>	FAB (MH <sup>+</sup> ) = 578
353						FAB (MH <sup>+</sup> ) = 592
354						FAB (MH <sup>+</sup> ) = 590




















































355					-CH <sub>3</sub>	m.p. = 233-235°C (dec)
356					-CH <sub>3</sub>	m.p. = 237-239°C (dec)
357					-CH <sub>3</sub>	m.p. = 228-229°C (dec)
358					-CH <sub>3</sub>	m.p. = 231-233°C (dec)
359					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 564
360					-CH <sub>3</sub>	m.p. = 225-226°C (dec)
361					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 571
362					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 629
363					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 645
364					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 670,668
365					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 571
366					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 585
367					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 632






















































368					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 613
369					-CH <sub>3</sub>	m.p. = 207-208°C (dec)
370					-CH <sub>3</sub>	m.p. = 229-220°C (dec)
371					-CH <sub>3</sub>	m.p. = 223-224°C (dec)
372					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 619
373					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 619
374					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 635
375					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 665
376					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 586
377					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 558
378					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 572
379					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 570
380					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 626
381					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 586

382			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 716
383			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	m.p. = 230-231°C (dec)
384			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	m.p. = 244-245°C (dec)
385			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 608
386			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 580
387			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 650
388			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 610
389			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 633
390			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 599
391			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	FAB(MH+) = 599
392			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	m.p. = 230-231°C (dec)
393			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	m.p. = 225-226°C (dec)
394			$\text{CH}_3$	$\text{CH}_3$	$-\text{CH}_3$	m.p. = 208-209°C (dec)

395					-CH <sub>3</sub>	m.p. = 218-219°C (dec)
396					-CH <sub>3</sub>	m.p. = 218-219°C (dec)
397					-CH <sub>3</sub>	m.p. = 229-230°C (dec)
398					-CH <sub>3</sub>	m.p. = 229-230°C (dec)
399					-CH <sub>3</sub>	m.p. = 215-216°C (dec)
400					-CH <sub>3</sub>	m.p. = 220-221°C (dec)
401					-CH <sub>3</sub>	m.p. = 224-225°C (dec)
402					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 673
403					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 663
404					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 596
405					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 671
406					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 608


407					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 594
408					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 582
409					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 640
410					-CH <sub>3</sub>	FAB(MH <sup>+</sup> ) = 619
411					-CH <sub>3</sub>	LRMS: calc'd:617 found (M+H <sup>+</sup> ): 618
412					-CH <sub>3</sub>	LRMS: calc'd:603 found (M+H <sup>+</sup> ): 604
413					-CH <sub>3</sub>	LRMS: calc'd:673 found (M+H <sup>+</sup> ): 674
414					-CH <sub>3</sub>	LRMS: calc'd:673 found (M+H <sup>+</sup> ): 674
415					-CH <sub>3</sub>	LRMS: calc'd:659 found (M+H <sup>+</sup> ): 660
416					-CH <sub>3</sub>	LRMS: calc'd:667 found (M+H <sup>+</sup> ): 668
417					-CH <sub>3</sub>	LRMS: calc'd:639 found (M+H <sup>+</sup> ): 640
418		-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CF <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:645 found (M+H <sup>+</sup> ): 646
419					-CH <sub>3</sub>	LRMS: calc'd:673 found (M+H <sup>+</sup> ): 674

420					-CH <sub>3</sub>	LRMS: calc'd:639 found (M+H <sup>+</sup> ): 640
421		-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:611 found (M+H <sup>+</sup> ): 612
422		-SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:597 found (M+H <sup>+</sup> ): 598
423					-CH <sub>3</sub>	LRMS: calc'd:623 found (M+H <sup>+</sup> ): 624
424					-CH <sub>3</sub>	LRMS: calc'd:637 found (M+H <sup>+</sup> ): 637
425		-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:547 found (M+H <sup>+</sup> ): 548
426		-SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:533 found (M+H <sup>+</sup> ): 534
427		-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	LRMS: calc'd:627 found (M+H <sup>+</sup> ): 628
428					-CH <sub>3</sub>	LRMS: calc'd:639 found (M+H <sup>+</sup> ): 640
429					-CH <sub>3</sub>	LRMS: calc'd:580 found (M+H <sup>+</sup> ): 581
430					-CH <sub>3</sub>	LRMS: calc'd:603 found (M+H <sup>+</sup> ): 604
431					-CH <sub>3</sub>	LRMS: calc'd:580 found (M+H <sup>+</sup> ): 581
432					-CH <sub>3</sub>	LRMS: calc'd:595 found (M+H <sup>+</sup> ): 596




433					-CH <sub>3</sub>	LRMS: calc'd:613 found (M+H <sup>+</sup> ): 614
434					-CH <sub>3</sub>	LRMS: calc'd:626 found (M+H <sup>+</sup> ): 627
435					-CH <sub>3</sub>	LRMS: calc'd:655 found (M+H <sup>+</sup> ): 656
528					-CH <sub>3</sub>	m.p. = 212-214°C (dec)
529		-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>			-CH <sub>3</sub>	m.p. = 215-217°C (dec)
530					-CH <sub>3</sub>	m.p. = 210-211°C (dec)
531					-CH <sub>3</sub>	m.p. = 223-225°C (dec)
532					-CH <sub>3</sub>	m.p. = 209-211°C (dec)
563			H	H		HRMS calc'd: 612.1935 found: 612.1952 m.p.=187-197°C
564			H	H		HRMS calc'd: 606.2638 found: 606.2638 m.p.=185-195°C
565			H	H		m.p. = 178-188°C
566			H	H		HRMS calc'd: 606.2638 found: 606.2633 m.p.=175-183°C
567			H	H		HRMS calc'd: 656.1430 found: 656.1438 m.p.=185-192°C
568						HRMS calc'd: 634.2951 found: 634.2968 m.p.=185-195°C

-147-

569		$-\text{SO}_2(\text{CH}_2)_2\text{CH}_3$	$\text{CH}_3$	$\text{CH}_3$		HRMS calc'd: 608.2464 found: 608.2477 m.p.=205-215°C
570		$-\text{SO}_2\text{CH}_2\text{CH}_3$	$\text{CH}_3$	$\text{CH}_3$		HRMS calc'd: 594.2308 found: 594.2311 m.p.=175-185°C
571			$\text{CH}_3$	$\text{CH}_3$		HRMS calc'd: 620.2794 found: 620.2805 m.p.=185-190°C




No.	W	R <sup>30</sup>	Physical Data
553	$-\text{SO}_2(\text{CH}_2)_2\text{CH}_3$	$-\text{CN}$	LRMS: calc'd: 573; found (M+1) <sup>+</sup> : 574
554		$-\text{CH}_3$	LRMS: calc'd: 574 found (M+1) <sup>+</sup> : 575
555		$-\text{CH}_3$	LRMS: calc'd: 576 found (M+1) <sup>+</sup> : 577
556	$-\text{SO}_2\text{CH}_2\text{CH}_3$	$-\text{CH}_3$	LRMS: calc'd: 548; found (M+1) <sup>+</sup> : 549
557	$-\text{SO}_2\text{CH}_2\text{CF}_3$	$-\text{CH}_3$	LRMS: calc'd: 602; found (M+1) <sup>+</sup> : 603
558		$-\text{CH}_3$	LRMS: calc'd: 560 found (M+1) <sup>+</sup> : 561



-148-

What is Claimed:



1. A compound having the structural formula




- 5 including all isomers and pharmaceutically acceptable salts, esters, and solvates thereof,

wherein one of Y and Z is N and the other is N, CH, or C-alkyl;

- X is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>6</sup>-, -CO-, -CH<sub>2</sub>-, -CS-, -C(OR<sup>5</sup>)<sub>2</sub>-,  
 -C(SR<sup>5</sup>)<sub>2</sub>-, -CONR<sup>20</sup>-, -C(alkyl)<sub>2</sub>-, -C(H)(alkyl)-, -NR<sup>20</sup>-SO<sub>2</sub>-,  
 10 -SO<sub>2</sub>-NR<sup>20</sup>-, -NR<sup>20</sup>CO-, -O-CO-NH-, -NH-CO-O-,
















R is







-149-

(wherein X<sup>1</sup> is -CH<sub>2</sub>-, -O-, or -NR<sup>7</sup>-),

10

-150-


- hydrogen, acyl, alkyl, alkoxy, alkenyl, cycloalkyl, cycloalkyl substituted with one or two groups selected from the group consisting of alkyl and carbonyl, cycloalkenyl, bicycloalkyl, arylalkenyl, benzyl, benzyl substituted with up to three independently selected  $R^3$  groups,
- 5 cycloalkylalkyl, polyhaloacyl, benzyloxyalkyl, hydroxy $C_2$ - $C_{20}$ alkyl, alkenylcarbonyl, alkylarylsulfonyl, alkoxycarbonylaminoacyl, alkylsulfonyl, or arylsulfonyl, additionally, when X is  $-CH_2-$ , R may also be  $-OH$ ; in further addition, when X is not N, R may also be hydroxymethyl, in further addition, R and X may combine to form the
- 10 group  $Prot-(NOAA)_r-NH-$  wherein r is an integer of 1 to 4, Prot is a nitrogen protecting group and when r is 1, NOAA is a naturally occurring amino acid or an enantiomer thereof, or when r is 2 to 4, each NOAA is a peptide of an independently selected naturally occurring amino acid or an enantiomer thereof;
- 15  $R^1$  and  $R^{21}$  are independently selected from the group consisting of H, alkyl, alkenyl, cycloalkyl, cycloalkenyl, bicycloalkyl, alkynyl, cyano, aminoalkyl, alkoxycarbonyl, aminocarbonyl, hydroxyamidino, alkoxycarbonylalkyl, phenyl alkyl, alkylcarbonyloxyalkyl,



H,  $-OH$ , (provided  $R^1$  and  $R^{21}$  are both not  $-OH$  and Y is not N), formyl,  $-CO$  alkyl,  $-CO$ acyl,  $-CO$ aryl, and hydroxyalkyl; additionally  $R^1$  and  $R^{21}$  together may form the group  $=CH_2$ ,  $=N-OR^5$ ,  $=N-CN$ ,  $=N-N(R^5)_2$ ,  $=CH-alkyl$ , alkylene,  $=C(alkyl)-alkyl$  or  $=C(halo)_2$ ; in further addition,  $R^1$


-151-

and R<sup>21</sup> together with the carbon atom to which they are attached may



or R<sup>1</sup> and R<sup>21</sup> together with the carbon atom to which they are attached may form a saturated heterocyclic ring containing 3 to 7 carbon atoms, one or more of which may be optionally substituted by alkyl, and one or two groups independently selected from S, O, and N-R<sup>20</sup>;

R<sup>2</sup> is



- R<sup>3</sup>, R<sup>4</sup>, R<sup>22</sup>, R<sup>24</sup>, and R<sup>25</sup> are independently selected from the group consisting of alkyl, H, halo, alkoxy, benzyloxy, benzyloxy substituted by nitro or aminoalkyl, haloalkyl, polyhaloalkyl, nitro, cyano, sulfonyl, hydroxy, amino, alkylamino, formyl, alkylthio, polyhaloalkoxy, acyloxy, trialkylsilyl, alkylsulfonyl, arylsulfonyl, acyl, alkoxycarbonyl alkylsulfinyl; -OCONH<sub>2</sub>, -OCONH-alkyl, alkylaminoalkyl, dialkylaminoalkyl, -COOH, -CON(R<sup>20</sup>)<sub>2</sub>, -OCON(alkyl)<sub>2</sub>, -NHCOO-alkyl, -NHCO-alkyl, phenyl, hydroxyalkyl, or morpholino;

- each R<sup>5</sup> and R<sup>6</sup> is independently selected from the group consisting of H and alkyl, provided that when X is C(OR<sup>5</sup>)<sub>2</sub> or C(SR<sup>5</sup>)<sub>2</sub>, both R<sup>5</sup> groups cannot be H, and in addition, when X is C(OR<sup>5</sup>)<sub>2</sub> or C(SR<sup>5</sup>)<sub>2</sub>, the two R<sup>5</sup> groups in X may be joined to form —(CR<sup>20</sup>)<sub>p</sub>— wherein p is an integer of 2 to 4;

R<sup>7</sup> is independently selected from the group consisting of H, alkyl, arylalkyl, cycloalkyl, aryl and aryl substituted with R<sup>3</sup> and R<sup>4</sup> as defined herein;

- each R<sup>8</sup> is independently selected from the group consisting of H, hydroxyalkyl or alkyl, or two R<sup>8</sup> groups may be joined to form an alkylene group;

R<sup>9</sup> is H, alkyl, aralkyl, or acyl;

R<sup>20</sup> is H, aryl or alkyl;

-152-

$R^{27}$  and  $R^{28}$  are independently selected from the group consisting of H, alkyl, hydroxyalkyl, arylalkyl, aminoalkyl, haloalkyl, thioalkyl, alkylthioalkyl, carboxyalkyl, imidazolylalkyl, and indolylalkyl; or  $R^{27}$  and  $R^{28}$  may combine to form an alkylene group;

5  $R^{29}$  is H, alkyl, -CO-alkyl, -CO-cycloalkyl, alkoxycarbonyl, aminocarbonyl, aryloxy carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfonyl, arylsulfonyl or -SO<sub>2</sub>-NH-R<sup>20</sup>;

$R^{30}$  is H, alkyl, aryl, cycloalkyl, hydroxyalkyl, aminoalkyl, -COOR<sup>20</sup>, -CON(R<sup>20</sup>)<sub>2</sub> or cyano;

10  $R^{31}$  and  $R^{32}$  are the same as  $R^{30}$  and in addition, two  $R^{30}$ ,  $R^{31}$  and  $R^{32}$  groups may form the group -(CH<sub>2</sub>)<sub>r</sub>- (wherein r is 1 to 6), in further addition,  $R^{31}$  and  $R^{32}$  can also be hydroxy, -N(R<sup>20</sup>)<sub>2</sub>, -O-acyl, -N(R<sup>20</sup>)acyl, -OCOOR<sup>20</sup>, or -OCON(R<sup>20</sup>)<sub>2</sub>;

15  $R^{33}$  is aryl or heteroaryl, with the proviso that when  $R^{33}$  is heteroaryl, the CO-R<sup>33</sup> bond is to a carbon atom in the  $R^{33}$  group;


$R^{34}$  is alkyl, cycloalkyl or aryl and in addition  $R^{34}$  may also be H when  $R^1$  and  $R^{21}$  together with the carbon atom to which they are attached form a saturated heterocyclic ring containing 3 to 7 carbon atoms and two groups independently selected from S, O, and N-R<sup>20</sup>;

20  $R^{35}$  is -CH<sub>2</sub>-, -NR<sup>20</sup>- or -O-;

$R^{36}$  is -NH<sub>2</sub>, alkyl or alkoxy;

$R^{37}$  is independently selected from the group consisting of H and alkyl;

$R^{38}$  is -CO-(CH<sub>2</sub>)<sub>0-5</sub>-OR<sup>5</sup>, -SO<sub>2</sub>-(alkyl), or



25 wherein  $q_1$  and  $q_2$  are independently 1-5, provided that the sum of  $q_1$  and  $q_2$  is 2-5; and

$R^{39}$  and  $R^{40}$  are independently selected from the group consisting of =O and (H,H).


30

2. A compound of claim 1 wherein Y and Z are N

3. A compound of claim 1 wherein Y is CH and Z is N


-153-

4. A compound of any of claims 1, 2 or 3 wherein R is




and X is O, SO, SO<sub>2</sub>, CH<sub>2</sub>, CH(alkyl), C(alkyl)<sub>2</sub>, CH(OH), or N(R<sup>20</sup>)CO.

5. A compound of any of claims 1, 2, 3, 4 or 5 wherein R<sup>3</sup> and R<sup>4</sup> are H; either R<sup>1</sup> is H, cycloalkyl or alkyl and R<sup>21</sup> is H or R<sup>1</sup> and R<sup>21</sup>


together form =O or  ; and wherein at least one of R<sup>27</sup> and R<sup>28</sup> is alkyl.

- 10 6. A compound of any of claims 1, 2, 3, 4 or 5 wherein R<sup>2</sup> is




and R<sup>30</sup> is H or CH<sub>3</sub>; R<sup>31</sup> and R<sup>32</sup> are H; and R<sup>33</sup> is ortho-substituted aryl or heteroaryl.

- 15 7. A compound of claim 6 wherein R<sup>33</sup> is



8. A compound of any one of claims 1 to 5 wherein R<sup>2</sup> is



- 20 and R<sup>34</sup> is methyl and R<sup>31</sup> and R<sup>32</sup> are H.

9. A compound as defined in claim 1 selected from the group consisting of compound numbers 17, 18, 25, 30, 31, 32, 34, 35, 36,

-154-

- 37, 41, 43, 44, 49, 53, 54, 56, 57, 58, 59, 80, 82, 84, 85, 94, 98, 100, 108, 121, 126, 127, 137, 145, 151, 152, 154, 155, 162, 166, 178, 179, 181, 185, 190, 191, 194, 199, 214, 215, 216, 225, 247, 253, 256, 257, 337, 339, 340, 341, 349, 351, 367, 409, 459, 479, 488, 489, 490, 500, 501, 502, 503, 505, 506, 507, 515, 516, 517, 555 and 562 from the tables of compounds appearing in the specification.
- 5
10. A pharmaceutical composition which comprises a compound as defined in claim 1 in combination with a pharmaceutically acceptable carrier.
- 10
11. A process for the preparation of a pharmaceutical composition as defined in claim 10 comprising admixing a compound as claimed in any of claims 1 to 9 with a pharmaceutically acceptable carrier.
- 15
12. The use of a compound of claim 1 for the preparation of a medicament for treating a cognitive or neurodegenerative disease.
- 20
13. A kit for treating a cognitive or neurodegenerative disease comprising in separate containers in a single package pharmaceutical compounds for use in combination, in one container a compound in accordance with claim 1 and in a separate container an acetylcholinesterase inhibitor, said compound and inhibitor each being in a pharmaceutically acceptable carrier and their combined quantities being an effective amount.
- 25
14. A method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a compound of claim 1, alone or in combination with an acetylcholinesterase inhibitor.
- 30



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :  
C07D 405/14, A61K 31/495  
C07D 409/14, 401/04, 413/14

A3

(11) International Publication Number: WO 98/05292

(43) International Publication Date: 12 February 1998 (12.02.98)

(21) International Application Number: PCT/US97/13383

(22) International Filing Date: 6 August 1997 (06.08.97)

(30) Priority Data:  
08/700,628 8 August 1996 (08.08.96) US

(71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US).

(72) Inventors: LOWE, Derek, B.; 237 North 9th Street, Kenilworth, NJ 07033 (US). CHANG, Wei, K.; 63 West Cedar Street, Livingston, NJ 07039 (US). KOZLOWSKI, Joseph, A.; P.O. Box 7391, Princeton, NJ 08543-7391 (US). BERGER, Joel, G.; 5675 North Camino Esplandora, Tucson, AZ 85718 (US). McQUADE, Robert; 221 Mountainview Avenue, Scotch Plains, NJ 07076 (US). BARNETT, Allen; 13 Flanders Drive, Pine Brook, NJ 07058 (US). SHERLOCK, Margaret; 34 Parkway West, Bloomfield, NJ 07003 (US). TOM, Wing; 133 Cedar Grove Parkway, Cedar Grove, NJ 07009 (US). DUGAR, Sundeeep; 749 Wingate Drive, Bridgewater, NJ 08807 (US). CHEN, Lian-Yong; 10 Pond View Drive, Plainsboro, NJ 08536 (US). CLADER, John, W.; 428 North Union Avenue, Cranford, NJ 07016 (US). CHACKALAMANNIL, Samuel; 79 Stratford Road, East Brunswick, NJ 08816 (US). WANG,

Yuguang; 38 Princess Drive, North Brunswick, NJ 08902 (US). McCOMBIE, Stuart, W.; 28 Hanford Place, Caldwell, NJ 07006 (US). TAGAT, Jayaram, R.; 133 Boynton Court, Westfield, NJ 07090 (US). VICE, Susan, F.; 1144 Sawmill Road, Mountainside, NJ 07092 (US). VACCARO, Wayne, D.; 1706 Westover Road, Yardley, PA 19067 (US). GREEN, Michael, J.; 1146 Rancho Encinitas Drive, Encinitas, CA 92024 (US). BROWNE, Margaret, E.; 26 Highfield Road, Bloomfield, NJ 07003 (US). ASBEROM, Theodoros; 525 Eagle Rock Avenue, West Orange, NJ 07052 (US). BOYLE, Craig, D.; 180 Northfield Road, Bridgewater, NJ 08807 (US). JOSTEN, Hubert, B.; 5441 Washington Boulevard, Jersey City, NJ 07310 (US).

(74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US).

(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


## Published

With international search report.

(88) Date of publication of the international search report:

2 April 1998 (02.04.98)

(54) Title: PIPERIDINE AND PIPRAZINE DERIVATIVES AND THEIR USE AS MUSCARINIC ANTAGONISTS



## (57) Abstract

Di-N-substituted piperazine or 1,4 di-substituted piperidine compounds in accordance with formula I (including all isomers, salts, esters, and solvates) wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>21</sup>, R<sup>27</sup>, R<sup>28</sup>, X, Y and Z as defined herein are muscarinic antagonists useful for treating cognitive disorders such as Alzheimer's disease. Pharmaceutical compositions and methods of preparation are also disclosed. Also disclosed are synergistic combinations of compounds of the above formula with acetylcholinesterase inhibitors.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



# INTERNATIONAL SEARCH REPORT

Int. Patent Application No.  
**PCT/US 97/13383**

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D405/14 A61K31/495 C07D409/14 C07D401/04 C07D413/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 96 26196 A (SCHERING CORP) 29 August 1996 see the whole document ---	1-13
A	EP 0 711 763 A (SANTEN PHARMA CO LTD) 15 May 1996 see claim 1 ---	1
A	US 5 010 078 A (ABOU-GHARBIA MAGID A ET AL) 23 April 1991 see claim 1 ---	1
A	EP 0 343 961 A (AMERICAN HOME PROD) 29 November 1989 see claim 1 ---	1
A	US 2 792 398 A (KYRIDES) 14 May 1957 see claim 1 ---	1
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

28 October 1997

Date of mailing of the international search report

13. 11. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 eponl,  
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

# INTERNATIONAL SEARCH REPORT

Int. National Application No.

PCT/US 97/13383

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CHEMICAL ABSTRACTS, vol. 45, no. 12,  25 June 1951  Columbus, Ohio, US;  abstract no. 5153,  OCHIACHI ET AL: "Electrolytic reduction  of the pyridinium ion."  column 1;  XP002044214  see abstract  &amp;  IBID,  pages 313-316,</p> <p style="text-align: center;">-----</p>	1

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/13383

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14  
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 14 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/13383

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9626196 A	29-08-96	AU 4971796 A	11-09-96
EP 0711763 A	15-05-96	FI 960364 A	26-01-96
		NO 960270 A	23-01-96
		CA 2168264 A	09-02-95
		CN 1128026 A	31-07-96
		WO 9504050 A	09-02-95
		JP 7089949 A	04-04-95
US 5010078 A	23-04-91	US 5482940 A	09-01-96
		US 5380725 A	10-01-95
		US 5106849 A	21-04-92
		US 5278160 A	11-01-94
		US 5254552 A	19-10-93
		AT 132862 T	15-01-96
		AU 628341 B	17-09-92
		AU 3502589 A	30-11-89
		DE 68925385 D	22-02-96
		DE 68925385 T	15-05-96
		DK 249989 A	25-11-89
		EP 0343961 A	29-11-89
		ES 2081302 T	01-03-96
		FI 94130 B	13-04-95
		GB 2218988 A,B	29-11-89
		IE 64151 B	12-07-95
		IL 90279 A	30-03-95
		JP 2015059 A	18-01-90
		PT 90633 B	30-11-94
EP 0343961 A	29-11-89	AT 132862 T	15-01-96
		AU 628341 B	17-09-92
		AU 3502589 A	30-11-89
		DE 68925385 D	22-02-96
		DE 68925385 T	15-05-96
		DK 249989 A	25-11-89
		ES 2081302 T	01-03-96
		FI 94130 B	13-04-95
		GB 2218988 A,B	29-11-89
		IE 64151 B	12-07-95
		IL 90279 A	30-03-95

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/13383

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0343961 A		JP 2015059 A	18-01-90
		PT 90633 B	30-11-94
		US 5482940 A	09-01-96
		US 5380725 A	10-01-95
		US 5010078 A	23-04-91
		US 5106849 A	21-04-92
		US 5278160 A	11-01-94
		US 5254552 A	19-10-93
-----			
US 2792398 A	14-05-57	NONE	
-----			

**THIS PAGE BLANK (USPTO)**